DRAFT TECHNICAL MEMORANDUM • FEBRUARY 2021

# Assessment of Groundwater Dependent Ecosystems for the Fillmore and Piru Basins Groundwater Sustainability Plan

PREPARED FOR

Fillmore and Piru Basins Groundwater Sustainability Agency P O Box 1110 Fillmore, CA 93016 PREPARED BY

Stillwater Sciences 2855 Telegraph Ave., Suite 400 Berkeley, CA 94705

# Suggested citation:

Stillwater Sciences. 2021. Assessment of Groundwater Dependent Ecosystems for the Fillmore and Piru Basins Groundwater Stability Plan. Technical Appendix. Prepared by Stillwater Sciences, Berkeley, California for Fillmore and Piru Basins Groundwater Sustainability Agency, Fillmore, California.

# **Table of Contents**

1	BACK	GROUND AND SETTING	1
	1.1	Physiography	1
	1.2	Aquifer and Soils	
	1.3	Hydrology	
	1.4	Historical Ecology	
2	GDE II	DENTIFICATION	8
	2.1	Vegetation Communities	
	2.1.		
	2.1.	_ 110000010	
	2.1.	1	
	2.2	Special-status Species	
	2.2.		
		Procedure	
	2.3	GDE units	16
3	GROU	NDWATER AND INTERCONNECTED SURFACE WATER HYDROLO	GY 19
	3.1	Groundwater Levels	19
	3.1.		
	3.1.	* *	
	3.2	Groundwater Quality	
		Interconnected Surface Waters	
	3.3.		
	3.3.		
	3.3.		
4	GDE C	ONDITION	30
	4.1	Esstadad Conditions	20
		Ecological Conditions	30
	4.1.	$\mathcal{E}$	
	4.1.		
	4.1.	- I I	
		Invasive Species	
		Ecological Value	69
	4.3.		
	4.3.	2 Fillmore	71
5	POTEN	ITIAL EFFECTS OF GROUNDWATER MANAGEMENT ON GDES	73
	5.1	Approach	73
	5.2	Biological Data	
	5.2.	~	
	5.2.		
	5.3	Climate Change Effects	
	5.4	Summary of Potential Effects	
		<b>→</b>	

	5.4.1 Piru 87	
	5.4.2 Fillmore	
	5.4.3 GDEs Important for GSP Analyses	0
6	GDE MONITORING10	1
7	PROJECTS AND MANAGEMENT ACTIONS10	1
8	LITERATURE CITED10	1
ΑF	ENDICES	1
ΑF	ENDIX A	2
ΑF	ENDIX B	3
ΑF	ENDIX C	5
Ta Ta Ta ass Ta	le 2.1-1. Vegetation and wetland data sources for Fillmore and Piru groundwater basins	6 7 0
Ta (N Ta oc Ta kn Ta	le 4.1-1. U.S. Fish and Wildlife Service (USFWS) and National Marine Fisheries Service (IFS) designated critical habitat <sup>1</sup> within the Fillmore and Piru groundwater basins	6.0
Ta the Ta kn Ta in	le 4.1-5. Special-status plant species and natural communities with known occurrence within Fillmore Groundwater Basin	3 7

Figures		
Figure 1.1-1.	Fillmore and Piru groundwater basins	3
Figure 1.4-1.	Riparian wetland forest extent along the Santa Clara River under historical and current conditions.	7
Figure 2.1-1.	Vegetation and wetland data sources used for the final GDE map	
Figure 2.1-2.	Final GDE determination based on the methods outlined in Section 2.1	
Figure 2.1-3.	Final GDE determination compared to iGDE mapping.	
Figure 2.3-1.	GDE units and groundwater wells for the Fillmore and Piru groundwater	17
1 iguic 2.5-1.	basins.	18
Figure 3.1-1.	Groundwater elevation and depth to groundwater measured at Well	10
11guic 3.1-1.	04N18W27B02S in the Del Valle Riparian Complex GDE Unit	20
Figure 2 1 2	Groundwater elevation and depth to groundwater measured at Well	20
Figure 3.1-2.	04N19W34K01S in the Santa Clara River Riparian Shrubland GDE Unit	21
Eigung 2 1 2	<u>-</u>	<i>∠</i> 1
Figure 3.1-3.	Groundwater elevation and depth to groundwater measured at Well	22
Fig. 2.1.4	04N19W34N01S in the Santa Clara River Riparian Shrubland GDE Unit	22
Figure 3.1-4.	Groundwater elevation and depth to groundwater measured at Well	22
E: 215	04N19W33D04S in the Cienega Riparian Complex GDE Unit	23
Figure 3.1-5.	Groundwater elevation and depth to groundwater measured at Well	~ .
TI 0.4.5	03N20W08A01S in the East Grove Riparian Complex GDE Unit	24
Figure 3.1-6.	Groundwater elevation and depth to groundwater measured at Well	
	03N20W09D01S in the East Grove Riparian Complex GDE Unit	25
Figure 3.3-1.	Dry season surface flow extent measured in the Fillmore Groundwater Basin	
	and the downstream portion of the Piru Groundwater Basin	29
Figure 4.1-1.	Five most common GDEs in the Del Valle Riparian Complex GDE Unit, by	
	acreage	31
Figure 4.1-2.	Five most common GDEs in the Santa Clara River Riparian Shrubland GDE	
	Unit, by acreage	31
Figure 4.1-3.	Five most common GDEs in the Cienega Riparian Complex GDE Unit, by	
	acreage	32
Figure 4.1-4.	Five most common GDEs in the Piru Creek Riparian Complex GDE Unit, by	
	acreage	32
Figure 4.1-5.	Five most common GDEs in the Tributary Riparian GDE Unit, by acreage	33
Figure 4.1-6.	Five most common GDEs in the East Grove Riparian Complex GDE Unit, by	
	acreage	34
Figure 4.1-7.	Five most common GDEs in the Sespe Creek Riparian Complex GDE Unit,	
	by acreage	35
Figure 4.1-8.	Critical habitat within the Fillmore and Piru groundwater basins	38
Figure 4.2-1.	Invasive species mapped within the Fillmore and Piru groundwater basins	68
Figure 5.2-1.	NDVI and NDMI for the Del Valle Riparian Complex GDE Unit in the Piru	
	Groundwater Basin.	75
Figure 5.2-2.	Slope of NDVI changes from 2011–2020 for the Del Valle Riparian	
C	Complex.	76
Figure 5.2-3.	NDVI and NDMI for the Santa Clara River Riparian Shrub GDE Unit in the	
C	Piru Groundwater Basin.	77
Figure 5.2-4.	NDVI and NDMI for the Piru Creek Riparian GDE Unit.	
Figure 5.2-5.	NDVI and NDMI for the Piru Basin Tributary Riparian GDE Unit	
Figure 5.2-6.	NDVI and NDMI for the Santa Clara River Riparian Shrub GDE Unit in the	. ,
	Fillmore Groundwater Basin.	80
Figure 5.2-7.	NDVI and NDMI for the Cienega Riparian Complex GDE Unit.	
1 15410 3.2 7.	1.2 .1 and 1.Divil for the Cichega Repartan Complex ODE Ont	01

Figure 5.2-8.	Slope of NDVI change in the Cienega Riparian Complex in the Fillmore and P.	iru
	Basins from 2011-2020. Red areas have declining NDVI and Blue areas have	
	increasing NDVI.	82
Figure 5.2-9.	NDVI and NDMI for the East Grove Riparian Complex GDE Unit	83
Figure 5.2-10	. Slope of NDVI change in the Cienega Riparian Complex in the Fillmore and	
_	Piru Basins from 2011-2020.	84
Figure 5.2-11	. NDVI and NDMI for the Fillmore Basin Tributary Riparian GDE Unit	85
Figure 5.2-12	NDVI and NDMI for the Sespe Creek Riparian Complex GDE Unit	86
Appendices		
	Vegetation Communities in the Fillmore and Piru Groundwater Basins	
	Special-status Terrestrial and Aquatic Wildlife Species from Database Queries w No Reliance on Fillmore or Piru Groundwater Dependent Ecosystem Units	ith
Appendix C.	Rooting Depths for Selected Species	

### 1 BACKGROUND AND SETTING

This Technical Appendix to the Fillmore and Piru Basins Groundwater Sustainability Plan (GSP) addresses the extent and condition of groundwater dependent ecosystems (GDEs) in the Fillmore and Piru Valley Groundwater Basins of the Santa Clara River Valley Groundwater Basin. The Fillmore and Piru groundwater basins are managed by Fillmore and Piru Basins Groundwater Sustainability Agency (FPBGSA). As part of the California's Sustainable Groundwater Management Act (SGMA), Groundwater Sustainability Agencies (GSAs) are required to consider GDEs and other beneficial uses of groundwater when developing their GSPs. SGMA defines GDEs as "ecological communities of species that depend on groundwater emerging from aquifers or on groundwater occurring near the ground surface" (23 CCR § 351(m)). As described in The Nature Conservancy's guidance for GDE analysis (Rohde et al. 2018), a GDE's dependence on groundwater refers to reliance of GDE species and/or ecological communities on groundwater for all or a portion of their water needs. Mapping GDEs requires mapping vegetation that can tap groundwater through their root systems, assessing where the depth of groundwater is within the rooting depth of that vegetation, and mapping the extent of surface water that is interconnected with groundwater (Rohde et al. 2018). Once the GDEs are mapped, the occurrence of specialstatus species can be used to assess the value of GDEs in the basin, while remote sensing measurements can be used to track the health of groundwater dependent vegetation through time. This information will be used to inform sustainable management criteria for each management unit. This appendix relies on hydrologic and geologic data presented in the GSP and its technical appendices.

### 1.1 Physiography

The Fillmore and Piru groundwater basins occupy the alluvial valley of the Santa Clara River from just west of the Los Angeles/Ventura County line to approximately 1 mile upstream of Santa Paula, California (Figure 1.1-1). The alluvial valley of the Santa Clara River serves as the floodplain of the Santa Clara River and the adjacent alluvial terraces and alluvial fans. The valley width ranges from approximately half a mile at the eastern boundary of the Piru subbasin, just west of the Los Angeles-Ventura county line, to over four miles at the confluence of Sespe Creek and the Santa Clara River. Together, the basins extend approximately 21 river miles along the valley axis and cover over 25,000 acres. Ground surface elevation ranges from 280 feet above sea level in the western Fillmore Groundwater Basin to about 1,000 feet above sea level in the eastern Piru Groundwater Basin and along the northern edge of the Fillmore Groundwater Basin (DWR 2016).

The basins are bounded to the north by Miocene to Pliocene marine deposits (the Pico Formation) of the Topatopa Mountains, uplifted by Tertiary thrusting along the San Cayetano Fault. The basins are bounded to the south by Oligocene continental deposits and Miocene to Pliocene marine deposits of the Oak Ridge and the Santa Susana mountains, uplifted by similar motion on the Oak Ridge Fault (Downs et al. 2020). The Santa Clara River occupies a valley between these ranges. Tectonic activity also influences drainage network patterns in the basins (Stillwater Sciences 2011).

Upstream of the Fillmore and Piru groundwater basins, the Santa Clara River is largely unregulated with the exception of Castaic Dam on Castaic Creek, and Bouquet Dam on Bouquet Creek. Castaic Dam regulates a large (398 km²) watershed and forms Castaic Lake, which is operated with Pyramid Lake as a hydroelectric pumped storage project. Because of the run-of-the-river operation of Castaic Dam, it unlikely to have a significant impact on Santa Clara River

low flow. Bouquet Dam regulates less than 1% of the Santa Clara River watershed area and is also unlikely to have a significant impact watershed hydrology (Stillwater Sciences 2011). There is a strong rainfall gradient from the relatively wet Sespe and Piru drainages in the western part of the watershed to the drier upper Santa Clara watershed in Los Angeles County,

The largest tributary to the Piru Groundwater Basin is Piru Creek, which joins the Santa Clara River about 6 miles downstream of the Piru Groundwater Basin boundary near Buckhorn. Piru Creek has a drainage area of 438 square miles and is blocked by Santa Felicia Dam and Lake Piru 5.7 miles upstream of the confluence with the Santa Clara River. The Piru Groundwater Basin extends up Piru Creek from the Santa Clara River to Santa Felicia Dam. Since construction of Santa Felicia Dam in 1955, flood flows on Piru Creek downstream of the dam have been regulated to below 990 cubic feet per second (cfs) except during the 1969 flood of record which had a peak flow of 28,800 cfs. Pyramid Dam was completed further upstream on Piru Creek in 1971 to impound water imported to the watershed via the California Water Project. Flows are released through Santa Felica Dam annually to recharge groundwater storage in the Fillmore and Piru groundwater basins, and other downstream basins, and to provide water for groundwater replenishment facilities further downstream. Hopper Canyon Creek joins the Santa Clara River approximately 3 miles downstream of the Piru Creek confluence.

The largest tributary in the Fillmore Groundwater Basin is Sespe Creek, which joins the Santa Clara River about 4.4 miles downstream of the boundary between the Fillmore and Piru groundwater basins just downstream of the town of Fillmore. Sespe Creek has a drainage area of 260 square miles and is largely unregulated. The older portions of the town of Fillmore were situated on the alluvial fan built by sediments from Pole Creek, a smaller (10.7-square mile) tributary between Sespe and Hopper Canyon creeks, which joins the Santa Clara River about 3 miles upstream of Sespe Creek. The drainage area of the Santa Clara River at the downstream end of the Fillmore Groundwater Basin is 1,476 square miles.




Figure 1.1-1. Fillmore and Piru groundwater basins.

### 1.2 Aguifer and Soils

The Fillmore and Piru groundwater basins are located in the distinctive geological province of the west-east trending Transverse Ranges. The Santa Clara River watershed has a drainage area of 1623 mi<sup>2</sup> that retains a relatively natural state compared with other large watersheds in coastal southern California. In general, both groundwater basins consist of an upper aquifer of sand and gravel Quaternary alluvium underlain by Pleistocene San Pedro Formation, which consists of permeable sand and gravels folded into an east-west trending syncline.

The Piru Groundwater Basin's eastern boundary coincides with exposures of shale and thinning of the surface alluvium to about 20 feet. The basin extends westward to the Fillmore Fish Hatchery, approximately one mile upstream of the City of Fillmore, where bedrock constriction of the alluvial basin results in increased groundwater elevation and seepage from the aquifer to the Santa Clara River. The most recent alluvium is typically 60-80 feet thick in the Piru Groundwater Basin, and is underlain by older alluvium, which occurs as terrace deposits up to 80 feet thick. Beneath the alluvium, the San Pedro Formation extends to depths of up to 8800 feet. Discontinuous clay layers also occur in the basin (UWCD 2016).

The Fillmore Groundwater Basin lies directly west of the Piru Narrows, a constriction in the valley width (Figure 1.1-1). The Fillmore Groundwater Basin includes the Santa Clara River alluvial valley, the Pole Creek fan, and Sespe Creek floodplain. The western boundary of the basin is located approximately half a mile east of the City of Santa Paula and coincides with high groundwater elevations and a gaining reach of the Santa Clara River. At the eastern boundary, recent alluvium is approximately 60 feet thick, deepening to 80 feet downstream. Older alluvium reaches depths of approximately 100 feet. Along the valley, the San Pedro formation reaches a depth of approximately 8400 feet; at the western boundary, it shallows to 5000-6000 feet. The northern portion of the basin is characterized by recent Quaternary alluvium. The most extensive alluvial fan, Pole Creek, is located between Sespe Creek and the Santa Clara River and underlies much of the City of Fillmore. West of Fillmore, the recent Sespe Creek alluvium is approximately 80 feet thick (UWCD 2016).

Soils in the basins are alluvial or eolian in origin and are derived from the upper Santa Clara River and surrounding mountains. Three soil associations occur in the basins (Edwards et al. 1970). The Santa Clara River floodplain is covered by Riverwash-Sandy Alluvial Land, an excessively drained, highly stratified complex of sediment ranging in size from sand to cobbles with minimal silt and clay. The Pico-Metz-Anacapa association occurs on alluvial terraces and extends into some of the westerly drainages in the Fillmore Groundwater Basin. This association is characterized by well- to excessively drained sandy loam and is 60 or more inches deep. Within this association, Anacapa soils are slightly alkaline. The Mocho-Sorrento-Garretson association occurs at higher elevations in tributaries at the mountain front, particularly on the north side of the basins. These well-drained loams to silty clay loams reach depths of over 60 inches and are some of the most agriculturally productive in Ventura County.

### 1.3 Hydrology

Natural surface flows in the Santa Clara River are supplemented by controlled releases from Lake Piru and, upstream of the Piru Groundwater Basin, releases from Castaic Lake. Major tributaries in the Fillmore and Piru groundwater basins include Piru, Hopper, and Sespe creeks. Apart from high flows associated with high-intensity rainfall events, flows in the Santa Clara River are generally low. At Montalvo, near the river mouth (USGS gage 11114000), daily flows are less

than 10 cfs for 50% of the year (Stillwater Sciences 2007d). Upstream of the Piru Groundwater Basin, at the county line (USGS gage 11108500), flow is typically 20-28 cfs (UWCD 2016). The low-flow hydrology of the Santa Clara River in the basin area has been characterized by alternating reaches of perennial and intermittent flow (Beller et al. 2016) based on bedrock constrictions and the resultant connection of the riverbed to the shallow aquifer. In the Fillmore and Piru groundwater basins, perennial flow occurs at the basin boundaries, with the extent of the perennial reaches varying based on inflows to the basin.

In the Fillmore and Piru groundwater basins, the Santa Clara River is a dynamic semi-confined braided stream with perennial reaches at the basin boundaries with large intermittent reaches between. The Santa Clara River Basin has an extremely high sediment supply and is subject to extremely large floods (Stillwater Sciences 2013). Minor channel shifting occurs during most 5–10-year floods (frequently associated with El Nino years), but the largest changes in channel alignment occur during less frequent larger magnitude floods that exceed a 10 year recurrence interval (Stillwater Sciences 2007, Stillwater Sciences 2013).

The shallow aquifers in both the Fillmore and Piru groundwater basins are considered to be unconfined aquifers with deeper aquifers having varying degrees of confinement. In the Piru Groundwater Basin, groundwater generally flows from east to west, parallel to the river channel, in both the alluvium and the San Pedro formation. At the bedrock constriction near the Fillmore Fish Hatchery, the narrower aquifer cross-section drives the water table upward, resulting in a perennial gaining reach of the Santa Clara River in all but the driest years. Downstream, in the Fillmore Groundwater Basin, the alluvium widens and groundwater continues to flow from east to west. Near the boundary of the Fillmore Groundwater Basin and the Santa Paula Basin to the west, low-permeability bedrock constricts the alluvium, albeit less severely than at the fish hatchery. Approaching the constriction, the narrower aquifer cross-section and southwest flow of recharged groundwater from the Sespe Creek alluvium drive the water table toward the surface. This reach of the Santa Clara River experiences perennial gaining conditions even in dry years. Minimum depth to groundwater was less than ten feet at both constrictions in 2015, during the drought (UWCD 2016)

The primary source of groundwater recharge is percolation from the Santa Clara River in the wider areas of the valley and alluvium, and from Piru and Sespe creeks. Groundwater outflow from the Piru Groundwater Basin to the east is also a significant source of recharge to the Fillmore Groundwater Basin. In both basins, recharge also occurs from stream percolation into outcrops of the San Pedro Formation to the north, from direct precipitation on the alluvium and northern alluvial fans, and from agricultural return flow.

Groundwater levels in both basins are sensitive to changes in surface flow. Annual conservation releases from Lake Piru by United Water resulted in approximately ten-foot increases in groundwater levels in the Piru Groundwater Basin and approximately five-foot increases in the Fillmore Groundwater Basin from 2009 to 2012 (UWCD 2016). Conversely, groundwater levels declined significantly during the 2013-2015 drought, during which no conservation releases were made.

### 1.4 Historical Ecology

The strongest process control on the distribution of floodplain wetland and riparian vegetation in the Santa Clara River is the availability of groundwater. Areas of rising groundwater that result in perennial surface flow and dense riparian woodlands are interspersed with 'losing reaches' with

intermittent surface flow conditions and less dense vegetation (Stillwater Sciences 2007a, Beller et al. 2011, Orr et al. 2011, Beller et al. 2016, Stillwater Sciences 2016). Within the forested wetland areas, flood disturbances are an important secondary control, with large floods causing significant adjustment of the channel and floodplain topography that periodically resets the pattern of vegetation. Surface erosion of the floodplain and bank erosion associated with lateral migration of channel causes uprooting of established trees while, subsequently, processes of deposition as the flood recedes established barren surfaces suitable for vegetation regeneration.

Together, surface water flows and fluvial geomorphic processes are the basis for vegetation recruitment and succession in GDEs, with surface water or shallow groundwater availability critical for vegetation growth. Such controls are common to GDEs, with previous studies of riparian vegetation in semi-arid river systems highlighting controls such as the magnitude and frequency of flood disturbance (Bendix 1994, 1997; Harris 1999; Bendix and Hupp 2000), depth to groundwater (as reflected in preference for gaining versus losing reaches; Stromberg et al. 1996, Shafroth et al. 1998), and a combination of the two (Hupp and Osterkamp 1996, Lite 2003, Bagstad et al. 2006, Leenhouts et al. 2006, Osterkamp and Hupp 2010).

Prior to Euro-American settlement in valley in the late 1700s, mature stands of riparian woodland or forest were focused on four locations in the Santa Clara River corridor. Evidence suggests that these locations had discrete patches of dense, persistent cottonwood-willow (*Populus* and *Salix* species) riparian forest, and that they corresponded to hydrologically gaining reaches of rising groundwater (Boughton et al. 2006, Stillwater Sciences 2007c, Beller et al. 2011, Orr et al. 2011, Beller et al. 2016). While the extent of these four riparian forests has reduced over time as population increased in the Santa Clara valley, their locations have been persistent, pointing to the fundamental role of hydrogeological control on forest existent (Beller et al., 2016). Three of the four locations—the East Grove, Cienega, and Del Valle—occur in the Fillmore and Piru groundwater basins (Figure 1.4-1), with the East Grove located at the downstream end of the Fillmore Groundwater Basin, Cienega at the boundary between the Fillmore and Piru groundwater basins, and Del Valle at the upstream end of the Piru Groundwater Basin (Figure 1.4-1). All are areas where groundwater occurs close to the floodplain surface.

Although the general locations suitable for forested wetlands are unchanged from historical conditions as a function of hydrogeology, surface water and shallow groundwater conditions are now influenced by surface water diversions, managed water releases, and treated wastewater releases. As a result, despite significant changes in general to the vegetation of the Santa Clara River valley and riparian corridor (see for example, descriptions presented in Beller et al. 2011 and Stillwater Sciences 2016), remnants of the four historically persistent riparian-forested wetlands that were dominated by willows and cottonwoods are still supported albeit in a more fragmented form (Orr et al. 2011, Beller et al. 2016). This decreasing extent mirrors changes in the Santa Clara valley in the Fillmore and Piru Basins. Between the mid-nineteenth century and 2005, the extent of active riparian corridor decreased by nearly half, from an estimated 15,500 acres to 8,000 acres (Beller et al., 2016). Levees and berms constructed to protect agriculture reduced the area of riparian forest relative to historical conditions, particularly in the Cienega and East Grove areas Downs et al. 2013, Beller et al. 2016).




Figure 1.4-1. Riparian wetland forest extent along the Santa Clara River under historical and current conditions. Note the persistence of the four main forested wetland patches identified from historical source from at least the early 1800's to the present. (Adapted from Beller et al. 2016 and Stillwater Sciences 2016).

### 2 GDE IDENTIFICATION

# 2.1 Vegetation Communities

Potential GDE units in the Fillmore and Piru groundwater basins were identified using the California Department of Water Resources' (DWR) indicators of groundwater dependent ecosystems (iGDE) database. The database, which is published online and referred to as the Natural Communities Commonly Associated with Groundwater dataset (DWR 2020), includes vegetation and wetland natural communities. These data were reviewed and augmented with additional vegetation mapping datasets to produce a map of final GDE units; additional information on vegetation community composition, aerial imagery, depth to groundwater modeled from local wells (where available), plant and species distributions in the area, and plant species rooting depths were also reviewed to support this determination. Maximum rooting depths from the literature are provided in Appendix C, Table C-1. Another way to explore the rooting depth of plants is to assess their elevation relative to the river channel surface (the relative elevation). Assuming that the groundwater elevation near the stream is similar to the stream elevation, we can assess the likely rooting depth of plants based on their relative elevation. Stillwater Sciences (2007a) assessed the relative elevation of various plant types in the Santa Clara River. Those results are provided in Appendix C, Table C-2.

### 2.1.1 Data sources

This section includes brief descriptions of the vegetation community data and other information sources used to identify and aggregate potential GDEs into final GDE units. The iGDE database (DWR 2020) was reviewed in a geographic information system (GIS) and used to generate a preliminary map to serve as the primary basis for initial identification of potential GDEs in the Fillmore and Piru groundwater basins. This dataset is a combination of the best available data obtained from the following publicly available sources:

- Vegetation Classification and Mapping Program (VegCAMP), California Department of Fish and Wildlife (CDFW)
  - o Santa Clara River Parkway Vegetation Database (Stillwater Sciences 2007b). *Imagery date: 2005; Minimum mapping unit (MMU): 1-acre.*
- Classification and Assessment with Landsat of Visible Ecological Groupings (CalVeg) –
   United States Department of Agriculture Forest Service (USDA 2014). *Imagery date:* 2005; *Minimum mapping unit (MMU):* 2.5-acre.
- National Wetlands Inventory Version 2.0 (NWI v2.0), U.S. Fish and Wildlife Service (USFWS 2018). *Minimum mapping unit (MMU): 0.5-acre*.

A more recent vegetation mapping effort for the Santa Clara River was completed in 2018. This dataset was used in the final GDE identification in place of the Santa Clara River Parkway Vegetation Database (Stillwater Sciences 2007b) referenced above.

 Vegetation Mapping of the Santa Clara River, Ventura County and Los Angeles County, California. Prepared for the Western Foundation of Vertebrate Zoology (Stillwater Sciences 2019). *Imagery date: 2016; Minimum mapping unit (MMU): 0.5-acre.*

The extent of the integrated data sources is shown in Table 2.1-1 and Figure 2.1-1.

Table 2.1-1. Vegetation and wetland data sources for Fillmore and Piru groundwater basins.

	Mapped area (acres)			
Data source	Fillmore	Piru	Total	
Vegetation	•	•	•	
CalVeg	17,997.0	6,489.6	24,486.6	
Vegetation Mapping of the Santa Clara River	4,580.4	4,405.6	8,986.0	
Wetland				
NWI	8.5	0.3	8.8	
Total <sup>1</sup>	22,585.9	10,895.4	33,481.4	

<sup>&</sup>lt;sup>1</sup> Totals may not appear to sum exactly due to rounding error.




Figure 2.1-1. Vegetation and wetland data sources used for the final GDE map.

### 2.1.2 Procedure

The steps for defining and mapping GDEs outlined in Rohde et al. (2018) were used as a guideline for this process. A decision tree was applied to determine when species or biological communities were considered groundwater dependent based on definitions found in SGMA (State of California 2014) and Rohde et al. (2018). This decision tree, created to systematically and consistently address the range of conditions encountered, is summarized below; the term 'unit' refers to an area with consistent vegetation and hydrology:

The unit is a GDE if groundwater is likely:

- 1. An important hydrologic input to the unit during some time of the year, AND
- 2. Important to survival and/or natural history of inhabiting species, AND
- 3. Associated with a regional ifer used as a regionally important source of groundwater

The unit is not a GDE if its hydrologic regime is primarily controlled by:

- 1. Surface discharge or drainage from an upslope human-made structure(s), surface as irrigation canal, irrigated fields, reservoir, cattle pond, or water treatment pond/facility.
- 2. Precipitation inputs directly to the unit surface. This excludes vernal pools from being GDEs where units are hydrologically supplied by direct precipitation and very local shallow subsurface flows from the immediately surrounding area.

Rohde et al. (2018) recommend that maps of likely GDEs be compared with local groundwater elevations to determine where groundwater is within the rooting depth of potential GDEs. Given uncertainties in extrapolating well measurements to GDEs and differences in surface elevation of wells and GDEs, Rohde et al. (2018) recommend assigning GDE status to vegetation communities either within 30 feet of the ground surface or where interconnected surface waters are observed. Groundwater elevation data for the spring of 2019 (United Water 2020) was converted to a depth to groundwater surface in GIS using LiDAR (USGS 2018). The groundwater elevation in spring 2019 was used because the groundwater elevation was higher (i.e., closer to the ground surface) than other available years.

The vegetation community mapping data sources identified in Section 2.1.1 were combined in GIS to create a groundwater basin-wide vegetation map. Consistent with Klausmeyer et al. (2018), the most recent and highest resolution mapping was prioritized over earlier and coarser scale datasets. The datasets were prioritized in the following order, with the highest priority data sources listed first: Vegetation Mapping of the Santa Clara River (Stillwater Sciences 2019) and CalVeg (USDA 2014). A crosswalk between the two mapping classification systems (i.e., Manual of California Vegetation [Vegetation Mapping of the Santa Clara River] and CalVeg) can be found in Appendix A (Table A-1).

Finally, additional wetland mapping was incorporated where vegetation data were coarse and did not accurately capture wetland features. These additional wetland data sources were incorporated unilaterally across the selected vegetation data source and were chosen to represent the best available data for the extent of each vegetation data source. CalVeg was supplemented with the iGDE wetland mapping (DWR 2020) which is derived from the National Wetlands Inventory (USFWS 2018). The Vegetation Mapping of the Santa Clara River dataset was mapped to a scale that did not require supplemental wetland data.

# 2.1.3 Refine potential GDE map

To inform the assessment of GDE condition and potential effects (Sections 3 and 4), the basin-wide vegetation and wetland map was reviewed and each community was assigned a groundwater dependence category (i.e., unlikely or likely; Figure 2.1-2). This determination was based on species composition and the groundwater dependency of dominant species, whether they were considered groundwater dependent by the iGDE database (DWR 2020), and wetland indicator status (Lichvar et al. 2016). Although Klausmeyer et al. (2018) includes species with upland facultative wetland indicator status (Lichvar et al. 2016) in their list of groundwater dependent mapping units, these upland facultative species were classified as unlikely to be groundwater dependent based on their typical elevation relative to the low flow channel in the Santa Clara River watershed. Figure 2.1-3 shows the differences between the iGDE database and the potential GDE map. Section 4.1 discusses the vegetation communities that were identified as likely to depend on groundwater.

These potential GDEs were overlaid with the depth to water dataset (as discussed in Section 2.1.2) in GIS and any communities occurring outside of 30-ft depth to groundwater were removed. The depth to water modeling data did not cover the entire extent of both the Fillmore and Piru groundwater basins; in these areas a manual review of potential GDEs was performed using aerial imagery. Coast live oak (*Quercus agrifolia*) communities were included as potential GDEs when they occurred in riparian areas or drainages; any mapped stands growing on a clear slope were excluded based on landscape position and improbable connection to groundwater.




Figure 2.1-2. Final GDE determination based on the methods outlined in Section 2.1.




Figure 2.1-3. Final GDE determination compared to iGDE mapping (Klausmeyer et al. 2018). The final GDEs are indicated by kept and added classes.

### 2.2 Special-status Species

As part of the ecological inventory, special-status species and sensitive natural communities that are potentially associated with GDEs in the Fillmore and Piru groundwater basins were identified. For the purposes of this document, special-status species are defined as those:

- listed, proposed, or under review as endangered or threatened under the federal Endangered Species Act (ESA) or the California Endangered Species Act (CESA);
- designated by CDFW as a Species of Special Concern;
- designated by CDFW as Fully Protected under the California Fish and Game Code (Sections 3511, 4700, 5050, and 5515);
- designated as Forest Service Sensitive according to the Regional Forester's Sensitive Species Management Guidelines listed per USFS Memorandum 2670 (USFS 2011);
- designated as Bureau of Land Management (BLM) sensitive;
- designated as rare under the California Native Plant Protection Act (CNPPA); and/or
- included on CDFW's most recent Special Vascular Plants, Bryophytes, and Lichens List (CDFW 2020a) with a California Rare Plant Rank (CRPR) of 1, 2, 3, or 4.

Sensitive natural communities are defined as vegetation communities identified as critically imperiled (S1), imperiled (S2), or vulnerable (S3) on the most recent California Sensitive Natural Communities List (CDFW 2020b).

### 2.2.1 Data Sources

Stillwater ecologists queried databases on regional and local occurrences and spatial distributions of special-status species within the Fillmore and Piru groundwater basins. Spatial database queries included potential GDEs plus a 1-mile buffer. Databases queried include:

- California Natural Diversity Database (CNDDB) (CDFW 2019);
- Vegetation Mapping of the Santa Clara River, Ventura County and Los Angeles County, California, prepared for the Western Foundation of Vertebrate Zoology (Stillwater Sciences 2019);
- California Native Plant Society (CNPS) Manual of California Vegetation (2020);
- eBird (2021); and
- The Nature Conservancy (TNC) freshwater species lists generated from the California Freshwater Species Database (CAFSD) (TNC 2020).

## 2.2.2 Procedure

Stillwater reviewed the database query results and identified special-status species and community types with the potential to occur within or be associated with the vegetation and aquatic communities in or immediately adjacent to the potential GDEs. Stillwater ecologists then consolidated these special-status species and sensitive community types into a list along with summaries of habitat preferences, potential groundwater dependence, and reports of any known occurrences (Section 4.1.31 and Appendix A).

Wildlife species were evaluated for potential groundwater dependence using determinations from the Critical Species Lookbook (Rohde et al. 2019) or by evaluating known habitat preferences, life histories, and diets. Species GDE associations were assigned one of three categories:

- Direct—species directly dependent on groundwater for some or all water needs (e.g., cottonwood with roots in groundwater, juvenile steelhead in dry season).
- Indirect—species dependent upon other species that rely on groundwater for some or all water needs (e.g., riparian birds).
- No known reliance on groundwater.

Sensitive natural communities were classified as either likely or unlikely to depend on groundwater based on species composition using the same methodology as vegetation communities (Section 2.1.3). Plant species were evaluated for potential groundwater dependence based on their habitat (Jepson Flora Project 2020) and association with vegetation communities classified as GDEs. Special-status plant GDE associations were assigned one of three categories: likely, possible, or unlikely. The 'possible' category was included to classify plant species with limited habitat data or where a species may have an association with a vegetation community identified as a GDE (e.g., Coast live oak, California sagebrush).

Database query results for local and regional special-status species occurrences were combined with their known habitat requirements to develop a list of groundwater dependent special-status species (Section 4.3.1) that satisfy the following criteria: (1) documented to occur within the GDE unit, or (2) known to occur in the region and suitable habitat present in the GDE unit.

### 2.3 GDE units

Seven GDE units were identified based on hydrologic and ecological conditions of the GDEs identified using the methods outlined in Section 2.1.3 (Table 2.3-1 and Figure 2.3-1). The exterior boundaries of each unit were established using the extent of GDE vegetation communities.

GDE units	Description				
Del Valle Riparian Complex	Historical Del Valle complex located at the upstream end of the Piru Groundwater Basin. Unit is predominantly dense riparian forest, with established Freemont cottonwood and red willow.				
Santa Clara River Riparian Shrubland	Riparian zone along the Santa Clara River; dominated by facultative phreatophytes and riparian shrubland habitat. Unit occupies both Fillmore and Piru groundwater basins. Unit is characterized by lower density and low stature shrubs and is dominated by mulefat.				
Cienega Riparian Complex	Historical Cienega complex located near the Fillmore Fish Hatchery. Unit occurs in both Fillmore and Piru groundwater basins. Unit is dominated by mulefat and giant reed of variable density throughout.				
East Grove Riparian Complex	Historical East Grove complex located at the downstream end of the Fillmore Groundwater Basin. Unit is occupied by dense riparian forest dominated by mulefat, black cottonwood and red willow.				
Piru Creek Riparian	Riparian zone along Piru Creek from Santa Felicia Dam to Highway 126. Unit is characterized by a thin but dense riparian corridor, dominated by mulefat, Freemont cottonwood and red willow.				
Tributary Riparian	Riparian habitat within tributaries to both Fillmore and Piru groundwater basins. Predominantly located to the north of the Santa Clara River draining the Topa Topa mountain range. Unit is dominated by oaks and other hardwoods.				
Sespe Creek Riparian	Riparian zone along Sespe Creek from the boundary of the Fillmore Groundwater Basin to Highway 126. Unit is dominated by mixed hardwood and low stature willows.				

Table 2.3-1. GDE unit description.

GDE units in the Fillmore and Piru groundwater basins contain 3,955.1 acres of GDE habitats, with the majority (65%) occurring in the Fillmore Groundwater Basin (Table 2.3-2). The Santa Clara River Riparian Shrubland and the East Grove Riparian complex are the largest GDE units and together account for 83% of the GDE units in the Fillmore Groundwater Basin.

Table 2.3-2. Extent of GDEs by GDE unit by Groundwater Basin (acres).

GDE unit	Fillmore	Piru	Total	
Cienega Riparian Complex	133.6	159.6	293.2	
Del Valle Riparian Complex	-	466.1	466.1	
Santa Clara River Riparian Shrubland	1,046.0	342.1	1,388.1	
East Grove Riparian Complex	1,101.9	-	1,101.9	
Piru Creek Riparian	-	336.9	336.9	
Tributary Riparian	196.6 68.9		265.5	
Sespe Creek Riparian	103.4	-	103.4	
Total	2,581.5	1,373.6	3,955.1	




Figure 2.3-1. GDE units and groundwater wells for the Fillmore and Piru groundwater basins.

# 3 GROUNDWATER AND INTERCONNECTED SURFACE WATER HYDROLOGY

This section uses groundwater level data to assess temporal trends in the groundwater elevations for GDE Unit with well data (Section 3.1). Section 3.2 explores groundwater quality and its potential effect on GDEs. Finally, Section 3.3 explores the extent of interconnected surface water groundwater quality and spatial trends in interconnected surface water that can later be used to assess potential impacts of groundwater management to GDEs.

### 3.1 Groundwater Levels

Historical dry periods and droughts play a major influence on groundwater elevations across the Fillmore and Piru groundwater basins. Droughts in 1974–1977, 1986–1991 and 2012–2016 have significant signatures in the hydrographs of shallow wells located besides the identified GDEs. Most recently, basins were near to full capacity in 2011 (i.e., groundwater levels very close to the surface) ahead of the 2012–2016 drought, which generally caused in water levels in wells to decline. The largest drops are seen in wells in the central part of Piru Groundwater Basin, and in the Sespe Upland and Pole Creek Fan area of Fillmore Groundwater Basin. The greatest seasonal fluctuations are seen in the same areas (central Piru Groundwater Basin, Sespe Upland, and Pole Creek Fan) where hydraulic gradients are relatively steep. In general, ground surface elevation in the Sespe Upland areas increases steeply to the north, unlike the relative flat topography near to the Santa Clara River channel. As such, depths to groundwater tend to be greatest in the Sespe upland area.

Long-term records of shallow groundwater are relatively rare in the Fillmore and Piru groundwater basins. Below we use well data presented in the Monitoring Plan and Data Gaps Analysis Appendix (DBS&A 2020) to explore shallow groundwater trends for the GDE Units shown in Figure 2.3-1 from east to west. We were unable to examine the groundwater levels in the Tributary Riparian GDE unit because there are no representative wells located in or near the unit. Many of the wells used in the analysis below are screened below the shallow groundwater depths used by GDEs (Table 3.1-1) and may not accurately represent the actual groundwater elevation. In addition, groundwater wells are absent in some of the GDE units (e.g., Tributary Riparian). The following sections analyze changes to groundwater elevations for some of the GDE units, because some of the wells aren't screened within the rooting zone of most GDEs, they may be indicative of changes in shallow groundwater levels through time, but may not represent the actual groundwater levels under the GDEs.

## 3.1.1 Del Valle Riparian Complex

Well 04N18W27B02S is an active agricultural well installed in 1932, with a screen depth of 140–255 feet (Table 3.1-1). The well is approximately 1500 ft north of the downstream end of the Del Valle Riparian Complex. The depth of groundwater in this well varies from about 19–160 feet below ground surface (bgs) (Figure 3.1-1). After a period of low water elevation levels in the 1960s, the water level between droughts was generally within 10-30 ft bgs. During droughts the water level drops to greater than 100 ft bgs, before recovering. The low water levels prior to the 1970s occur before the effluent releases from Los Angeles County increased. Groundwater elevation have increased since the 2012-2016 drought, but by early 2020 had not yet reached the long-term average depth (Figure 3.1-1).

**Table 3.1-1.** Characteristics for wells used for the groundwater quality and groundwater level assessment. The location of the wells is shown on Figure 2.3-1.

Well	Basin	GDE Unit	Screen depth (ft bgs)	Data type <sup>1</sup> Water quality data available		Water level data available	
04N18W27B02S	Piru	Del Valle Riparian Complex	140–255	WQ, WL	1960–2004	1932–2020	
04N19W34K01S	Piru	Santa Clara River Riparian Shrubland	5–120	WQ, WL	1931–2016	1972–2019	
04N19W33H01S	Piru	Cienega Riparian Complex	237–362	WQ	1954–1958	2007–2011**	
04N19W33D04S	Fillmore	Cienega Riparian Complex	140–486	WQ, WL	1951–1985	1972–2020	
03N20W03N01S	Fillmore	Santa Clara River Riparian Shrubland	120–172	WQ, WL	1981–1994	1959–2020	
03N20W09D01S	Fillmore	East Grove Riparian Complex	210–310	WQ, WL	1969–2019	1988–2019	
03N20W08A01S	Fillmore	East Grove Riparian Complex	0-n/a	WQ, WL	1962–1969**	1959–2020	

<sup>&</sup>lt;sup>1</sup> WQ=groundwater quality, WL= groundwater level




Figure 3.1-1. Groundwater elevation and depth to groundwater measured at Well 04N18W27B02S in the Del Valle Riparian Complex GDE Unit.

# 3.1.2 Santa Clara River Valley Riparian Shrubland

Well 04N19W34K01S within the Santa Clara River Valley Riparian Shrubland GDE Unit in the Piru Basin and 0.45 miles east of the Cienega Riparian Complex boundary, has a screen depth of 5–120 feet bgs and was installed in 1972. The average depth to groundwater over the period or record was 19.5 feet bgs Groundwater levels in well declined sharply to 49-62 ft bgs during droughts (Figure 3.1-2), but quickly recovered to the long-term average values. Since 2016, water elevations have risen, but still have not fully recovered to a normal range of 15 feet bgs or less by fall 2019.



Figure 3.1-2. Groundwater elevation and depth to groundwater measured at Well 04N19W34K01S in the Santa Clara River Riparian Shrubland GDE Unit.

Well 03N20W03N01S is in the Fillmore Groundwater Basin, 0.1 miles outside of the Santa Clara River Valley Riparian Shrubland boundary (0.75 miles east of the East Grove Riparian Complex) and has been operational since 1959. This well has a screen depth of 120–172 feet bgs (Table 3.1-1). Over the 60 years of recorded depths to groundwater, this well averages 14.7 feet bgs (Figure 3.1-3). Similar to the other well, since 1970, the water elevation drops during droughts, with the largest drop occurring in 2016, when the water elevation was greater than 40 ft bgs. Subsequent to the droughts the water elevation recovers to its average long-term value.



Figure 3.1-3. Groundwater elevation and depth to groundwater measured at Well 04N19W34N01S in the Santa Clara River Riparian Shrubland GDE Unit.

### Cienega Riparian Complex

Well 04N19W33D04S lies within the Cienega Riparian Complex in the Fillmore Groundwater Basin and is screened at 140–486 feet (Table 3.1-1). In became operational in 1972 and the average depth to water from 1972–2019 was 5.4 feet bgs (Figure 3.1-4). The well data show significant declines during droughts, with groundwater level dropping to 25–33 feet during the droughts before returning to 5-10 ft bgs between the droughts.




Figure 3.1-4. Groundwater elevation and depth to groundwater measured at Well 04N19W33D04S in the Cienega Riparian Complex GDE Unit.

### East Grove Riparian Complex

Well 03N20W08A01S has data from 1959–2020, however the screen depth has not been recorded and has a high frequency of water level fluctuations not observed in the other wells (Figure 3.1-5). The average depth to water over the period of record is 13 feet bgs. Similar to the other wells, the water level declines during droughts, with the lowest water level of approximately 30 ft bgs in 2016. The recovery following the 2012–2016 drought was slower than previous droughts, but as of late 2019 was less than 10 ft bgs.




Figure 3.1-5. Groundwater elevation and depth to groundwater measured at Well 03N20W08A01S in the East Grove Riparian Complex GDE Unit.

Well 03N20W09D01S, in operation since 1988, is an agricultural well that has a screen depth of 210–310 feet bgs (Table 3.1-1). The average water level from 1988–2019 was 8.7 feet bgs (Figure 3.1-6). Groundwater data are relatively sparse in this well, particularly after 2010. This well shows a similar pattern to other wells described above with declines in groundwater elevation during droughts followed by recovery. During the 2012–2016 drought, recorded groundwater depths decreased to almost 25 ft bgs. Adam Lambert (UCSB) has established numerous groundwater monitoring points (<12 ft) in the East Grove Riparian Complex. These wells show similar results to Figure 3.2-5, but the magnitude of changes varies based on the location and relative elevation.




Figure 3.1-6. Groundwater elevation and depth to groundwater measured at Well 03N20W09D01S in the East Grove Riparian Complex GDE Unit.

### 3.2 Groundwater Quality

In general, groundwater quality in the Fillmore and Piru groundwater basins is high with a few exceptions (DBS&A 2020). Between 1951 and 1968, elevated concentrations of TDS, sulfate, chloride and boron near the Ventura/Los Angeles County Line were generally connected to loosely regulated oil fields in Los Angeles County. Some of these elevated concentrations persisted after the Clean Water Act reduced impacts from the oil fields (DBS&A 2020). High chloride concentrations in the Santa Clara River originating in Los Angeles County cause the largest concern for water quality, particularly for agricultural uses in the Piru Groundwater Basin. Historically water quality chemicals of concern (COCs) in the basins include:

- Total dissolved solids (TDS)
- Sulfate
- Chloride
- Nitrate
- Boron

The inter-quartile range of concentration for the five contaminants of concern and the EPA maximum contaminant level (MCL) in drinking water are shown in Table 3.2-1. In general, while concentrations for the contaminants of concern exceeded MCLs at times, they were not sufficiently high to impact GDEs. Moreover, shallow groundwater (used by GDEs) is usually younger and recharged by the surface waters of the Santa Clara watershed and has less legacy contamination than deeper groundwater (VCWPD 2016).

Table 3.2-1. Inter-quartile range and peak concentration (in square brackets) for selected contaminants observed in wells in the GDE units.

Well number	Basin	GDE unit	TDS (mg/l)	Chloride (mg/l)	Sulfate (mg/l)	Nitrate (mg/l)	Boron (mg/l)
04N18W27B02S	Piru	Del Valle Riparian Complex	2413–2696 [3123]	150–151 [159]	365–1412 [1536]	14–44 [66]	0.7–0.73 [1]
04N19W34K01S	Piru	Santa Clara River Riparian Shrubland	1366–1520 [1520]	41–55 [67]	616–339 [710]	9.2–13.6 [28]	0.6–0.8 [1.0]
04N19W33D04S	Piru	Cienega Riparian Complex	966–1180 [1221]	31 to 47 [49]	458-522 [1332]	9.8–16 [32]	1–1.1 [1.1]
04N19W33H01S	Fillmore	Cienega Riparian Complex	980 [n/a]	33–54 [54]	487–720 [720]	8.5–42 [42]	0.86–0.89 [0.89]
03N20W03N01S	Fillmore	Santa Clara River Riparian Shrubland	1048–1300 [1400]	41–62 [70]	476–580 [580]	17–41 [48]	0.8–0.9 [0.9]
03N20W09D01S	Fillmore	East Grove Riparian Complex	1290–1516 [1516]	74–79 [81]	587–616 [650]	39.5–73 [73]	0.5–0.8 [0.9]
MCL (mg/l)			500 <sup>a</sup>	250 <sup>a</sup>	250 <sup>a</sup>	45 <sup>b</sup>	1 <sup>c</sup>

 <sup>&</sup>lt;sup>a</sup> EPA secondary Maximum Contaminant Level
 <sup>b</sup> EPA Maximum Contaminant Level

<sup>&</sup>lt;sup>c</sup> California State Notification Level

### 3.3 Interconnected Surface Waters

Surface waters within the Piru and Fillmore groundwater basins have varying degrees of connection to groundwater. Generally, surface waters can be categorized into three major surface water types: perennial reaches, intermittent reaches, and ephemeral reaches. For this discussion we define perennial reaches as those that have surface flow except during prolonged and severe droughts. Surface flows occur for prolonged periods in intermittent reaches, but the reaches tend to go dry during most years. Ephemeral reaches occur in smaller tributaries within the basin and only support surface flow immediately after storm events.

As discussed in Section 1.4, the Santa Clara River has alternating perennial and intermittent reaches with perennial reaches occurring where rising groundwater contributes the vast majority of the surface water (except during storm events with significant runoff) and the intermittent reaches are losing reaches that are disconnected from groundwater during most of the year. Perennial reaches include the Del Valle Riparian Complex, the Cienega Riparian Complex, and the East Grove Riparian Complex (Figure 1.4-1). Some portion of the perennial reaches are downstream of the connected reach and the water has not completely infiltrated into the bed. Continuous surface water flow across the Fillmore and Piru basins can also be episodically established through effluent from Los Angeles County or dam releases such as the reach immediately downstream of Piru Dam.

### 3.3.1 Piru Groundwater Basin

The Piru Groundwater Basin has three perennial reaches that generally co-occur with the Del Valle Riparian Complex, the Cienega Riparian Complex, and Piru Creek from Santa Felicia Dam to the confluence with the Santa Clara River (Table 3.4-1). The Del Valle Reach occurs at the upstream end of the Piru Groundwater Basin where the valley width is still narrow and the alluvial basin is shallow relative to downstream reaches. The Del Valle Reach was historically perennial (Beller et al. 2016) and flows are currently augmented by effluent releases in Los Angeles County. The extent of dry season surface flow in the Cienega Reach was mapped from 2011–2018 by United Water and is discussed below. Several small ephemeral tributaries to the Santa Clara River and Piru Creek occur in the reach and are disconnected from groundwater. To our knowledge, there has not been a systematic exploration of the extent of surface water in lower Piru Creek.

Flows in the Del Valle Reach are monitored at the Santa Clara River near Piru gage (USGS Gage Number 11109000) from 1928–1932 and since water year 1997 at the La Falda Way Bridge and at the USGS gage at the Ventura County Line (USGS Gage Number 11108500) from 1953–1996. These gage records show that this reach is perennial, but shallow groundwater levels are not monitored in this reach. The degree to which interconnected surface waters are maintained by effluent releases in Los Angeles County is not known, however, are believed to be a significant contributor to surface water flow.

A 6.5-mile-long intermittent, losing reach occurs between Del Valle and the Cienega Reach of the Santa Clara River in the Piru Groundwater Basin. The groundwater is relatively deep in this reach and when the Santa Clara River flows, the surface flow is disconnected from groundwater. During prolonged surface water releases from Santa Felicia Dam or Castaic Lake, for example, the high stream bed infiltration rates will replenish the groundwater beneath and near the river and can cause groundwater levels to reach the stream bed. This condition will abate when the

surface water flows cease (or decrease dramatically) and the mounded groundwater beneath the river declines and the surface water flow returns to a disconnected state with the groundwater.

Flow in Piru Creek is maintained by releases from Santa Felicia Dam which likely also raises the groundwater level in this area. This reach of Piru Creek was historically intermittent (Beller et al. 2016) but there is currently perennial flow at least in the upper portion of the reach.

### 3.3.2 Fillmore Groundwater Basin

The Santa Clara River in the Fillmore Groundwater Basin has two perennial reaches that correspond to the Cienega Riparian Complex and the East Grove Riparian Complex. Both these reaches are typically dominated by rising groundwater except during storm flows. The rising groundwater flows can be supplemented by man-made releases from Santa Felicia Dam or Castaic Lake. The extent of rising groundwater (the source of the perennial flow) varies based on water year type (see section 3.3.3 for a discussion). Between these two reaches the Santa Clara River is an intermittent, losing reach for approximately 5 miles.

Sepse Creek is perennial at the USGS gage near the Fillmore Groundwater Basin boundary (where the creek exits the mountains) but the downstream extent of perennial flow is not known, although the lower portion of Sespe Creek is intermittent. Other tributaries within the Fillmore Groundwater Basin, including Pole Creek, Boulder Creek, and Timber Creek are typically ephemeral or intermittent. United Water mapped the extent of surface water from 2011-2017 and that is explored below.

#### 3.3.3 Variations in the extent of surface water 2011-2017

United Water used field observations and photographs to map the extent of dry-season surface water in the Santa Clara River from 2011–2017 (Figure 3.3-1). This period includes relatively wet 2011 and the 2012–2016 drought. The wetted area was mapped as extending downstream to the Freeman Diversion (red triangle in Figure 3.3-1) if a diversion occurred according to the diversion rate operations log. Santa Clara River flow rates are measured at the Fillmore Fish Hatchery (eastern circle in Figure 3.3-1, downstream end of Piru Groundwater Basin) and Willard Road (the western circle in Figure 3.3-1, downstream end of Fillmore Groundwater Basin). Sespe Creek flow is measured at the USGS gage (northernmost circle in Figure 1, August 2011 and June 2012). Castaic Lake and Santa Felicia Dam releases were absent for six months prior to the measurements shown on Figure 3.3-1 and the flows are therefore assumed to be entirely groundwater dependent.

Figure 3.3-1 shows that the extent of surface water and the magnitude of surface flows both decreased during the drought. The total length of the wetted stream channel in the Ventura County portion of the Santa Clara River decreased from approximately 17 miles in Fall 2012 to less than five miles in Fall 2016 (Figure 1). Streamflow at the Fillmore Fish Hatchery declined from 25 cfs in 2011 to 6 cfs in fall 2013. Surface flows were absent near the fish hatchery from 2014 to 2016 before returning in 2017. Streamflow at the downstream end of the Fillmore Groundwater Basin declined from 42 cfs in August 2011 to 1 cfs in October 2016. Subsequent dry-season flow measurements at Willard Road in 2018 and 2019 (not shown) ranged up to 16 cfs. Lower Sespe Creek has remained dry from Spring 2013 through November 2017. Surface flows at both the fish hatchery and Willard Road are linked with groundwater elevation.




Figure 3.3-1. Dry season surface flow extent measured in the Fillmore Groundwater Basin and the downstream portion of the Piru Groundwater Basin. The solid colored lines correspond to the extent of surface flow with the flow in cfs indicated in bubbles. The red triangle is the Freeman Diversion Dam, downstream of the Fillmore and Piru groundwater basins. Reaches where the downstream extent of surface water flow are inferred based on diversion at Freeman Dam are indicated by dotted lines. Figure courtesy of United Water.

### 4 GDE CONDITION

This section characterizes the GDE units based on their hydrologic and ecological conditions and assigns a relative ecological value to each unit by evaluating its ecological assets and its vulnerability to changes in groundwater (Rohde et al. 2018).

### 4.1 Ecological Conditions

GDEs included terrestrial and aquatic habitat and other open water aquatic habitats. There are few shallow groundwater wells in the Fillmore and Piru groundwater basins, but many of the deeper wells show that there continues to be shallow groundwater and interconnected surface water at the basin boundaries at the historical Del Valle, Cienega, and East Grove riparian woodlands (Figure 1.4-1). The Cienega and East Grove riparian woodlands are located at sites of rising groundwater due to constrictions in the valley width. The role of shallow groundwater elsewhere in the basin is less certain and will be assessed based on interpolated groundwater elevation and vegetation.

The GDE determination (i.e., likely or unlikely) is shown in Figure 2.1-2.

### 4.1.1 Vegetation Communities and GDE habitats

#### Piru

The Piru Groundwater Basin contains 1,373 acres of mapped GDEs which compose 13% of the total area of the basin. The Del Valle Riparian Complex contains the most GDEs within the Piru Groundwater Basin along with the Santa Clara River Riparian Shrubland and Piru Creek Riparian GDE units. GDEs within the basin are predominately riparian communities. The most prevalent vegetation community across all GDE units is mulefat thickets which makes up 41% of all mapped GDEs in the basin. Red willow riparian woodland and forest and Freemont cottonwood woodland and forest (*Populus fremontii* Forest Alliance) make up 19% and 17% of the GDEs basin-wide, respectively.

The Del Valle Riparian Complex GDE Unit contains 35% (458.1 acres) of mapped GDE units within the basin and is predominantly riparian habitat. Freemont cottonwood woodland and forest (127.7 acres), and red willow riparian woodland and forest (120.8 acres) are the dominant GDE types within this unit, with mulefat thickets (73.7 acres) also present throughout (Figure 4.1-1). These dominant vegetation communities are associated with the riparian zone on the mainstem Santa Clara River The maximum rooting depths reported in the literature for the cottonwood and willows is 6.9 ft, while the Mulefat (*Baccharis salicifolia*) has maximum reported rooting depths of about 2 ft (Appendix C). The depth to groundwater at monitoring well 04N18W27B02S is just less than 30 ft, but this well is located downstream of the Del Valle Riparian Complex GDE Unit, and likely records deeper groundwater than what occurs under the GDE.




Figure 4.1-1. Five most common GDEs in the Del Valle Riparian Complex GDE Unit, by acreage.

The Santa Clara River Riparian Shrubland GDE Unit contains 25% (329.8 acres) of mapped GDE units within the basin and is predominantly riparian habitat. Mulefat thickets (267.4 acres) is the dominant GDE type within this unit, with common and giant reed marshes (50.3 acres) also present throughout (Figure 4.1-2). These herbaceous and shrub communities, which are associated with the riparian zone of the mainstem Santa Clara River, are more tolerant of the drier conditions in this unit.




Figure 4.1-2. Five most common GDEs in the Santa Clara River Riparian Shrubland GDE Unit, by acreage.

The Cienega Riparian Complex GDE Unit contains 12% (159.6 acres) of the mapped GDE units within the basin and is predominantly riparian habitat. Mulefat thickets (84.7 acres) and red willow riparian woodland and forest (44.0 acres) and are the dominant GDE types within this unit (Figure 4.1-3). These dominant vegetation communities are associated with the riparian zone and the historical Cienega wetland complex on the mainstem Santa Clara River. Prior to the 2012–2016 drought, cottonwoods and willows were much more common in this unit. Similar to the East Grove Riparian Complex the maximum rooting depths range from 2 ft for mulefat to 6.9 ft for the cottonwood and willow forests.




Figure 4.1-3. Five most common GDEs in the Cienega Riparian Complex GDE Unit, by acreage.

The Piru Creek Riparian GDE Unit contains 24% (314.9 acres) of the mapped GDE units within the basin and is predominantly riparian habitat. Mulefat thickets (118.9 acres), Freemont cottonwood woodland and forest (267.4 acres) is the dominant GDE type within this unit, (92.8 acres), and red willow riparian woodland and forest (81.8 acres) are the dominant GDE types within this unit (Figure 4.1-4). These dominant vegetation communities are associated with the riparian zone of Piru Creek, which is a tributary to the Santa Clara River. Similar to the East Grove Riparian Complex the maximum rooting depths range from 2 ft for mulefat to 6.9 ft for the cottonwood and willow forests. Perennial flow in Piru Creek may sustain some of the plants.




Figure 4.1-4. Five most common GDEs in the Piru Creek Riparian Complex GDE Unit, by acreage.

The Tributary Riparian GDE Unit contains 4% (58.3 acres) of the mapped GDE units within the basin and is predominantly composed of Riparian Mixed Shrub Alliance (26.0 acres) and Baccharis (Riparian) Alliance (18.5 acres) (Figure 4.1-5). These vegetation communities are associated with drainages to Piru Creek and the Santa Clara River, including Holser and Hopper Canyons. Vegetation along Holser Canyon is primarily comprised of Riparian Mixed Shrub Alliance, while the vegetation along Hopper Canyon is a mixture of Riparian Mixed Shrub Alliance and Riparian Mixed Hardwood. Maximum rooting depths for these vegetation types aren't well documented but likely range from 2.0 ft for *Baccharis* species (i.e., Riparian Mixed Scrub Alliance) to 6.9 ft or deeper for the cottonwood and willow species present in Riparian Mixed Hardwood Alliance. This GDE unit contains no shallow groundwater data to assess groundwater linkages, but the vegetation is likely not tied to the aquifer.




Figure 4.1-5. Five most common GDEs in the Tributary Riparian GDE Unit, by acreage.

## **Fillmore**

The Fillmore Groundwater Basin contains 2,582 acres (acres) of mapped GDEs which compose 11% of the total area of the basin. The East Grove Riparian Complex and Santa Clara River Riparian Shrubland GDE units compose the majority of the GDEs within the Fillmore Groundwater Basin and GDEs within the basin are predominately riparian communities. The most prevalent vegetation community across all GDE units is mulefat thickets (*Baccharis salicifolia* Shrubland Alliance) which makes up 47% of all mapped GDEs in the basin. Red willow riparian woodland and forest (*Salix laevigata* Woodland Alliance) and black cottonwood woodland and forest (*Populus trichocarpa* Forest Alliance) make up 14% and 13% of the GDEs basin-wide, respectively.

The East Grove Riparian Complex GDE Unit contains 43% (1,101.9 acres) of mapped GDE units within the basin and is predominantly riparian habitat. Mulefat thickets (357.5 acres), red willow riparian woodland and forest (317.0 acres), and black cottonwood woodland and forest (315.5 acres) are the dominant GDE types within this unit (Figure 4.1-6). These dominant vegetation communities are associated with the riparian zone and the historical East Grove wetland complex on the mainstem Santa Clara River. Similar to the Del Valle Riparian Complex, the maximum rooting depths range from 2 ft for mulefat to 6.9 ft for the cottonwood and willow forests. The

groundwater elevation in the East Grove Riparian Complex is within the rooting depth of the vegetation types that make up the GDE unit during most years.




Figure 4.1-6. Five most common GDEs in the East Grove Riparian Complex GDE Unit, by acreage.

The Santa Clara River Riparian Shrubland GDE Unit contains 40% (1,046 acres) of mapped GDE units within the basin and is also predominantly riparian habitat. Mulefat thickets (743.6 acres) and the common and giant reed marshes association (*Phragmites australis - Arundo donax* Herbaceous Semi-Natural Alliance; 134.3 acres) are the dominant GDE types within this unit (Figure 4.1-2). These herbaceous and shrub communities, which are associated with the riparian zone of the mainstem Santa Clara River, are more tolerant of the drier conditions in this unit. Mulefat (*Baccharis salicifolia*) has maximum reported rooting depths of about 2 ft and arundo, the dominant plant species of the common and giant reed marshes Herbaceous Semi Natural Alliance has rooting depths up to 16 ft (Appendix C). Comparing the rooting depths to the range of groundwater depths in Figure 3.2-2 shows that groundwater elevations are often within the maximum root depth of Arundo and may be within the rooting depth of mulefat at lower relative elevations during particularly wet years, although this was not common.

The Cienega Riparian Complex GDE Unit contains 5% (133.6 acres) of the mapped GDE units within the basin and is predominantly riparian habitat. Common and giant reed marshes (64.8 acres) and mulefat thickets (59.0 acres) are the dominant GDE types within this unit (Figure 4.1-3. These dominant vegetation communities are associated with the riparian zone and the historical Cienega wetland complex on the mainstem Santa Clara River. The rooting depth ranges from 2 ft for mulefat to 16 ft for arundo (Appendix C). Mulefat thickets typically occur at higher relative elevation in the Santa Clara River despite their shallow rooting depth suggesting that they are often disconnected from groundwater. Prior to the 2012-2016 drought Fremont cottonwoods and willows (both with maximum rooting depths of 6.9 ft were common. Kibler et al. (2019) have shown that much of the cottonwood and willow species died out during the drought where the drop in groundwater was 16.4 ft or greater. The mean relative elevation of willows in the Santa Clara River ranges from 4.8 to 12.4 ft, while Fremont cottonwoods have mean relative elevation of 9.7 ft (Appendix C, Table C–2). Common and giant reed (Arundo) have a maximum reported rooting depth of 16 ft and occur at a mean relative elevation of 7.6 ft (Appendix C).

The Tributary Riparian GDE Unit contains 8% (196.6 acres) of the mapped GDE units within the basin and is predominantly composed of Coast Live Oak Alliance (193.4 acres) with Riparian Mixed Hardwood Alliance also present (3.2 acres) (Figure 4.1-5). These vegetation communities are associated with drainages from the mountain range to the north of the basin (i.e., Santa Paula Ridge and San Cayetano Mountain). Coast Live Oak has rooting depth up to 35.1 ft, while the Riparian Mixed Hardwood Alliance has rooting depths up to 6.9 ft (Appendix C). Neither the groundwater depth nor the source of water for vegetation in this GDE is known. The tributaries are not perennial which suggests that a combination of surface water and groundwater may support the riparian hardwood. The Coast Live Oak alliance may tap other water sources.

The Sespe Creek Riparian Complex GDE Unit contains 4% (103.4 acres) of the mapped GDE units within the basin and is predominantly riparian habitat. Willow (shrub) Alliance (19.6 acres) and Riparian Mixed Hardwood Alliance (19.2 acres) are the dominant GDE types within this unit (Figure 4.1-7). These dominant vegetation communities are associated with the riparian zone of Sespe Creek, which is a tributary to the Santa Clara River. Maximum rooting depths for both willows and cottonwoods in the literature is 6.9 ft (Appendix C, Table C–2). The mean relative elevation of willows range from 4.8 to 12.4 ft in the Santa Clara River, while Fremont cottonwoods have mean relative elevation of 9.7 ft (Appendix C, Table C–2). Data on the depth to groundwater in sparse but can exceed 30 ft in places.




Figure 4.1-7. Five most common GDEs in the Sespe Creek Riparian Complex GDE Unit, by acreage.

### 4.1.2 Beneficial uses

The Water Quality Control Plan (Basin Plan) for the Los Angeles Region (LARWQCB 2014) identifies the surface waters in the GDE units as having a variety of beneficial uses pertaining to fish, wildlife, and GDEs. These beneficial uses apply to aquatic features that are fed by groundwater within the Fillmore and Piru groundwater basins. The beneficial uses for aquatic features and groundwater vary between aquatic features and include:

• Groundwater recharge (GWR);

- Freshwater replenishment (FRSH);
- Warm freshwater habitat (WARM);
- Cold freshwater habitat (COLD);
- Wildlife habitat (WILD);
- Preservation of biological habitats of special significance (BIOL);
- Support of habitat for rare, threatened, or endangered species (RARE);
- Warm and cold migration habitat (MIGR);
- Warmwater spawning habitat (SPWN); and
- Wetland habitat (WET).
- Aquaculture (AQUA).

Beneficial uses include those that directly benefit groundwater conditions (e.g., groundwater recharge [GWR]), those supported directly by groundwater via interconnected surface waters (e.g., freshwater replenishment [FRSH]; support of rare, threatened, or endangered species [e.g., Southern California steelhead, California condor] [RARE]), and those that apply to groundwater beneficial uses (i.e., aquaculture [AQUA]).

# 4.1.3 Special-status species

The Fillmore and Piru groundwater basins are ecologically important and provide habitat for numerous wildlife species that are groundwater dependent. Within the two groundwater basins, five plants, 11 natural communities, 10 wildlife, and four fish species were identified as indirectly or directly groundwater dependent and may occur within the Fillmore and Piru groundwater basins. Appendix B provides information for special-status terrestrial and aquatic wildlife species from the database queries that are not groundwater dependent and/or unlikely to occur in the GDE units, including each species' regulatory status, habitat associations, and documented occurrences in the groundwater basins.

The Fillmore and Piru groundwater basins include designated critical habitat for four federally listed species: California condor (*Gymnogyps californianus*), Least Bell's vireo (*Vireo bellii pusillus*), southwestern willow flycatcher (*Empidonax traillii extimus*), and Southern California steelhead (*Oncorhynchus mykiss*) (USFWS 1976, USFWS 1977, USFWS 1994, USFWS 2013, NMFS 2005). The amount of critical habitat for each species within the Fillmore and Piru groundwater basins is shown in Table 4.1-1 and locations are shown in Figure 4.1-8.

Table 4.1-1. U.S. Fish and Wildlife Service (USFWS) and National Marine Fisheries Service (NMFS) designated critical habitat<sup>1</sup> within the Fillmore and Piru groundwater basins.

Common name	USFWS critical habitat (acres)			NMFS critical habitat (miles)		
Scientific name	Fillmore	Piru	Total	Fillmore	Piru	Total
California condor Gymnogyps californianus	2	1	2	-	1	-
Least Bell's vireo Vireo bellii pusillus	-	1,443	1,443	-	1	-
Southwestern willow flycatcher Empidonax traillii extimus	2,472	2,612	5,083	-	-	-
Southern California steelhead Oncorhynchus mykiss	-	-	-	15.4	15.3	30.7

All species	2,474	4,055	6,528	15.4	15.3	30.7

<sup>&</sup>lt;sup>1</sup> Data sources: USFWS 1976, USFWS 1977, USFWS 1994, USFWS 2013, NMFS 2005

Habitat management and special-status species recovery plans have been implemented in the Fillmore and Piru groundwater basins and include protections for special-status species and associated habitats. These plans include *United Water Conservation District Multiple Species Habitat Conservation Plan* (UWCD 2018), *Southern California Gas Company Multi-Species Habitat Conservation Plan* (SoCalGas 2020), *Santa Clara River Upper Watershed Conservation Plan* (TNC 2006), *Conservation Plan for the Lower Santa Clara River Watersheds and Surrounding Areas* (TNC 2008), and *Santa Clara River Enhancement and Management Plan* (VCWPD and LADPW 2005).




Figure 4.1-8. Critical habitat within the Fillmore and Piru groundwater basins.

# Piru

### Plants and natural communities

Four potentially groundwater dependent special-status plant species were documented in the Piru Groundwater Basin (Table 4.1-2). One species, white rabbit-tobacco, was identified as likely to depend on groundwater and has been observed in the open rock and sand wash bed of the Santa Clara River within the Del Valle Riparian Complex and Cienega Riparian Complex GDE units. Three species were identified as possibly dependent on groundwater and are predominantly associated with the Tributary Riparian GDE Unit. Slender mariposa lily (*Calochortus clavatus* var. *gracilis*) and Greata's aster are both associated with Foothill Canyon habitats and San Fernando Valley spineflower (*Chorizanthe parryi* var. *fernandina*), which has been observed directly upstream of the Piru Groundwater Basin, is associated with coastal sage scrub habitat.

Ten potentially groundwater dependent sensitive natural communities were documented with the potential to occur in the Piru Groundwater Basin (Table 4.1-2). Seven of these communities are identified as likely to depend on groundwater; these GDEs are predominantly riparian habitats (e.g., Southern cottonwood willow riparian forest, Southern willow scrub and California sycamore woodlands [*Platanus racemosa – Quercus agrifolia* Woodland Alliance]) and occur throughout the basin in all of the GDE units present within the Piru Groundwater Basin (i.e., Del Valle Riparian Complex, Santa Clara River Riparian Shrubland, Cienega Riparian Complex, Piru Creek Riparian, and Tributary Riparian). Additional potentially groundwater dependent sensitive natural communities include arrow weed thickets [*Pluchea sericea* Shrubland Alliance], which is a shrubland communities associated with intermittent or seasonally flooded washes and is documented in the Del Valle Riparian Complex.

Table 4.1-2. Special-status plant species and sensitive natural communities with known occurrence within the Piru Groundwater Basin.

Common name Scientific name	Status <sup>1</sup>	Association with GDE	Occurrence location	Source <sup>2</sup>	Habitat and occurrence
Plants					
Slender mariposa lily Calochortus clavatus var. gracilis	1B.2, S2S3, G4T2T3, not state or federally listed	Possible	Del Valle Riparian Complex, Piru Creek Riparian, Tributary Riparian	CNDDB	Shaded foothill canyons, chaparral; eight regional CNDDB observations, primarily in sagebrush on north-facing slopes and ridges of Newhall Ranch area.
San Fernando Valley spineflower Chorizanthe parryi var. fernandina	1B.1, S1, G2T1, SE, FPT	Possible	Tributary Riparian	CNDDB	Sandy coastal scrub, valley and foothill grassland; single CNDDB record from 2011 in Potrero Canyon near its confluence with the Santa Clara River.
Payne's bush lupine Lupinus paynei	1B.1, S1, G1Q, not state or federally listed	Unlikely	Piru Creek Riparian	CNDDB	Coastal and riparian scrub, valley and foothill grassland, generally on sandy soils; single CNDDB record from 2008 in Rancho Temescal.
Ojai navarretia Navarretia ojaiensis	1B.1, S2, G2, not state or federally listed	Unlikely	Del Valle Riparian Complex, Tributary Riparian	CNDDB	Openings in chaparral or coastal scrub, valley and foothill grassland, generally on clay soils; four CNDDB records on clay soil in grasslands of Ventura Homestead and Newhall Ranch areas.
White rabbit-tobacco Pseudognaphalium leucocephalum	2B.2, S2, G4, not state or federally listed	Likely	Cienega Riparian Complex, Del Valle Riparian Complex	CNDDB	Sandy or gravelly benches, dry stream or canyon bottoms; CNDDB records in sandy, rocky washes.
Greata's aster Symphyotrichum greatae	1B.3, S2, G2, not state or federally listed	Possible	Tributary Riparian	CNDDB	Mesic areas in broadleafed upland forest, chaparral, cismontane and riparian woodland; CNDDB records in Hopper and Pine canyons.

Common name Scientific name	Status <sup>1</sup>	Association with GDE	Occurrence location	Source <sup>2</sup>	Habitat and occurrence
Sensitive Natural Commu	nities				
Arrow weed thickets  Pluchea sericea  shrubland alliance	S3, G4	Likely	Del Valle Riparian Complex	VMSCR	Around springs, seeps, irrigation ditches, canyon bottoms, stream borders, and seasonally flooded washes; occur in the Central Valley and desert areas of southern California.
California sycamore woodlands Platanus racemosa – Quercus agrifolia woodland alliance	S3, G3	Likely	Piru Creek Riparian	VMSCR	Riparian woodlands along intermittent streams, springs, seeps, riverbanks, and floodplain terraces; occur in scattered stands along California's central and southern Coast Ranges, the Sierra Nevada foothills, the Peninsular and Transverse ranges, and the western Mojave and Colorado deserts.
California walnut groves  Juglans californica  forest and woodland  alliance	\$3, G3	Unlikely	Piru Creek Riparian, Tributary Riparian	CalVeg, CNDDB, VMSCR	Riparian corridors, but most stands cover all hillslopes; occur along southern California coast and in Peninsular and Transverse ranges.
Fremont cottonwood forest and woodland Populus fremontii – Fraxinus velutina – Salix gooddingii forest and woodland alliance	S3, G4	Likely	Del Valle Riparian Complex, Piru Creek Riparian, Santa Clara River Riparian Shrubland	CalVeg, VMSCR	On floodplains, along low-gradient rivers and streams, and in alluvial fans and valleys with a dependable subsurface water supply; occur throughout much of California except the Sierra Nevada and Modoc Plateau.
Scale broom scrub  Lepidospartum  squamatum shrubland alliance	\$3, G3	Unlikely	Piru Creek Riparian, Santa Clara River Riparian Shrubland	CalVeg, VMSCR	Intermittently or rarely flooded, low-gradient alluvial deposits along streams, washes, and fans; occur in southern California in the inner central Coast Ranges, Transverse and Peninsular ranges, and the Mojave Desert.
Southern cottonwood willow riparian forest	S3.2, G3	Likely	Piru Creek Riparian	CNDDB	Frequently flooded lands along rivers and streams; occur in the Transverse and Peninsular ranges from Santa Barbara County south and east to the edge of the deserts.
Southern mixed riparian forest	S2.1, G2	Likely	Tributary Riparian	CNDDB	
Southern riparian scrub	S3.2, G3	Likely	Santa Clara River Riparian Shrubland	CNDDB	Sand and gravel bars close to river channels; occur along and at the mouths of perennial and intermittent streams of the southern Coast Ranges.

Common name Scientific name	Status <sup>1</sup>	Association with GDE	Occurrence location	Source <sup>2</sup>	Habitat and occurrence
Southern willow scrub	S2.1, G3	Likely	Del Valle Riparian Complex, Piru Creek Riparian, Santa Clara River Riparian Shrubland	CNDDB	Generally on alluvium deposited near stream channels during floods; occur along major rivers of coastal southern California
Valley oak riparian	S2.1, G3	Unlikely	Tributary Riparian	CNDDB	Valley bottoms and gentle slopes that are intermittently flooded; occur in the Coast Ranges, Central Valley, and foothills of the Sierra Nevada, Cascade, and Klamath ranges.

<sup>1</sup> Status codes:

G = Global

= Subspecies or variety

Federal

FT = Listed as threatened under the federal Endangered Species Act

FPT = Proposed as threatened under the federal Endangered Species Act

FD = Federally delisted

Rank Critically Imperiled—At very high risk of extinction due to extreme rarity (often 5 or fewer populations), very steep declines, or other factors. 1

State

- Imperiled—At high risk of extinction due to very restricted range, very few populations (often 20 or fewer), steep declines, or other factors.
- Vulnerable At moderate risk of extinction or elimination due to a restricted range, relatively few populations (often 80 or fewer), recent and widespread declines, or other factors.

S = Sensitive

SSC = CDFW species of special concern

SFP = CDFW fully protected species

SE = Listed as Endangered under the California Endangered Species Act ST = Listed as Threatened under the California Endangered Species Act

- Apparently Secure Uncommon but not rare; some cause for long-term concern due to declines or other factors.
- Demonstrably Secure Common; widespread and abundant.
- Taxonomic questions associated with this name

Ranks such as S2S3 indicate a ranking between S2 and S3

#### California Rare Plant Rank (CRPR)

- 1B Plants rare, threatened, or endangered in California and elsewhere
- 2B Plants rare, threatened, or endangered in California, but more common elsewhere
- More information needed about this plant, a review list
- 4 Plants of limited distribution, a watch list

#### CRPR Threat Ranks:

- 0.1 Seriously threatened in California (high degree/immediacy of threat)
- 0.2 Fairly threatened in California (moderate degree/immediacy of threat)
- 0.3 Not very threatened in California (low degree/immediacy of threats or no current threats known)
- <sup>2</sup> Sources: CNDDB (CDFW 2019), CalVeg (USDA 2014), VMSCR (Vegetation Map of the Santa Clara River; Stillwater Sciences 2019).

# Terrestrial and aquatic wildlife

Ten potentially groundwater dependent special-status terrestrial and aquatic wildlife species were identified as having the potential to occur within the Piru Groundwater Basin: one amphibian species, seven bird species, and two reptile species (Table 4.1-3). Additional information on these species, including regulatory status, habitat associations, and documented occurrences in the groundwater basin, is provided in Table 4.1-3.

Southwestern pond turtle and two-striped garter snake are the only wildlife species likely present (i.e., documented occurrences) in the Piru Groundwater Basin classified as directly groundwater dependent due to their association with riverine and lentic habitats. Southwestern pond turtles likely use habitat within the Del Valle Riparian Complex, Cienega Riparian Complex, Piru Creek Riparian GDE units for foraging (e.g., open water), nesting (e.g., grasses and forbs alliance), and/or overwintering (e.g., mixed riparian and willow scrub and cottonwood or willow alliance). Two-striped garter snake likely use freshwater (e.g., perennial riverine) and riparian (e.g., riparian mixed hardwood and/or shrub) habitats within the Delle Valle Riparian Complex, Cienega Riparian Complex, and Piru Creek Riparian GDE units for capturing prey. One special-status, Arroyo toad (*Anaxyrus californicus*), was not documented in the Piru Groundwater Basin but is included species could possibly occur along the Santa Clara River and within the Delle Valle Riparian Complex, Cienega Riparian Complex, and Piru Creek Riparian GDE units (VCWPD and LADPW 2005, Stillwater Sciences 2007c).

Indirectly groundwater dependent bird species use riparian habitat (e.g., willow/willow shrub, cottonwood, mixed riparian alliances) within the Del Valle Riparian Complex, Santa Clara River Riparian Shrubland, Cienega Riparian Complex, and Piru Creek Riparian GDE units for foraging, nesting, and migratory habitat. These GDE units include designated critical habitat for Southwestern willow flycatcher (Del Valle Riparian Complex [433 acres], Santa Clara River Riparian Shrubland [316 acres], Cienega Riparian Complex [154 acres], and Piru Creek Riparian [246 acres]), and Least Bell's vireo (Del Valle Riparian Complex [436 acres] and Tributary Riparian [6 acres]).

Table 4.1-3. Groundwater-dependent special-status terrestrial and aquatic wildlife species with known occurrence or suitable habit in the Piru Groundwater Basin.

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in Piru Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Piru Groundwater Basin
Amphibian			,		_	
Arroyo toad Anaxyrus californicus	FE/SSC	Possible (suitable habitat)	No documented occurrences	CAFSD	Direct	Washes, arroyos, sandy riverbanks, riparian areas with willows, sycamores, oaks, cottonwoods; needs exposed sandy streamsides with stable terraces for burrowing, with scattered vegetation for shelter, and areas of quiet water or pools free of predatory fishes with sandy or gravel bottoms without silt for breeding. Arroyo toads depend on groundwater for breeding in shallow still pools and riparian vegetation that provides foraging habitat (Rohde et al. 2019). Historically found in the upper and lower Santa Clara River basin and currently persists in Middle/Upper Piru Creek upstream of Lake Piru (Santa Clara River Trustee Council 2008). A habitat assessment conducted by Stillwater Sciences (2007c) estimated that there were 893 acres of potential habitat along the Piru River and Santa Clara River upstream of the confluence with Piru Creek.
Reptile	•		•	•	•	•
Two-striped gartersnake Thamnophis hammondii	BLMS, FSS/SSC	Likely	Cienega Riparian Complex, Delle Valle Riparian Complex, Piru Creek Riparian	CNDDB, CAFSD	Direct	Highly aquatic snake species. Found in or near permanent fresh water, often along streams with rocky beds and riparian vegetation. Prey includes fish, fish eggs, tadpoles, newt larvae, small frogs and toads, leeches, and earthworms. Commonly found in the Santa Clara River Watershed (UWCD 2018). Occurrences on Santa Clara River between Salt Creek and Summer Four Crossing in 2000; in spillway ponds downstream of Lake Piru in 2016, along Piru Creek near Piru Canyon Road in 2009 (CDFW 2019).

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in Piru Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Piru Groundwater Basin
Southwestern pond turtle Actinemys pallida <sup>5</sup>	BLMS, FSS/SSC	Likely	Cienega Riparian Complex, Del Valle Riparian Complex, Piru Creek Riparian	CNDDB, CAFSD	Direct	Ponds, lakes, rivers, streams, creeks, marshes, and irrigation ditches with basking sites. Feeds on aquatic plants, invertebrates, worms, frog and salamander eggs and larvae, crayfish, and occasionally frogs and fish. Relies on surface water that may be supported by groundwater (Rhode et al. 2019). Documented throughout the Santa Clara River and in parts of the Piru Creek watersheds. Occurrences include Santa Clara River upstream of Piru in 2000, spillway ponds and downstream of Lake Piru in 2016 (CDFW 2019).
Bira	1		I	l	I	D
California condor Gymnogyps californianus	FE/SE	Likely	Piru Creek Riparian	CNDDB	Indirect	Require vast expanses of open savannah, grasslands, and foothill chaparral in mountain ranges of moderate altitude; deep canyons containing clefts in rocky walls and large trees provide nest sites; forages up to 100 miles from roost to nest. Forage in grasslands, oak savanna habitats, and Condors may rely on groundwater dependent vegetation for nesting in foothill grasslands, oak savanna habitats, and oldgrowth forest (Rohde et al. 2019). Condors observed drinking water at a small perched pool near spillway canyon just west of the Santa Felicia Dam in the vicinity of the Piru Creek Riparian GDE Unit and within Piru Canyon (CDFW 2019, eBird 2021).

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in Piru Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Piru Groundwater Basin
Least Bell's vireo Vireo bellii pusillus	FE/SE	Likely	Cienega Riparian Complex, Del Valle Riparian Complex, East Grove Riparian Complex, Piru Creek Riparian, Sespe Creek Riparian	CNDDB, CAFSD	Indirect	Nests in dense vegetative cover of riparian areas; often nests in willow or mulefat; forages in dense, stratified canopy. This species relies on groundwater-dependent vegetation in riparian areas, particularly during breeding periods (Rohde et al. 2019). Eats insects, fruits, and berries. Documented throughout the Santa Clara River and Piru Creek in the Piru Groundwater Basin (CDFW 2019, eBird 2021, WFVZ 2020c). Documented breeding occurs on the TNC Sespe Cienega property (Bruce citation, eBird 2021). Critical habitat located along the Santa Clara River in the Del Valle Riparian Complex GDE Unit (USFWS 1994).
Southwestern willow flycatcher Empidonax traillii extimus	FE/SE	Likely	Delle Valle Riparian Complex	CNDDB, CAFSD	Indirect	Dense brushy thickets within riparian woodland often dominated by willows and/or alder, near permanent standing water. Reliant on groundwater-dependent riparian vegetation, including for nest sites that are typically located near slow-moving streams, or side channels and marshes with standing water and/or wet soils (Rohde et al. 2019). Feeds on insects, fruits, and berries. Occurrences throughout the Santa Clara River (CDFW 2019, eBird 2021). Critical habitat located along the Santa Clara River in the Del Valle Riparian Complex and Piru Creek GDE units between the confluence with Santa Clara River and Lake Piru (USFWS 2013).
Western yellow- billed cuckoo Coccyzus americanus occidentalis	BLMS, FSS, FT/SE	Likely	Del Valle Riparian Complex	CNDDB	Indirect	Summer resident of valley foothill and desert riparian habitats; nests in open woodland with clearings and low, dense, scrubby vegetation. Reliant on groundwater-dependent riparian vegetation for habitat (Rhode et al. 2019). Historical populations documented along the Santa Clara River 3.5 miles east of Piru in 1979 are possibly extirpated (CDFW 2019).

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in Piru Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Piru Groundwater Basin
White-tailed kite Elanus leucurus	BLMS/SFP	Likely	Cienega Riparian Complex, Del Valle Riparian Complex, Piru Creek Riparian, Santa Clara River Riparian Shrubland	CNDDB	Indirect	Lowland grasslands and wetlands with open areas; nests in trees near open foraging area. Predominately preys on small mammals, but its diet also includes birds, and lizards. Occurrences along Santa Clara River and Piru Creek (CDFW 2019, eBird 2021, WFVZ 2020c).
Yellow warbler Setophaga petechia	-/SSC	Likely	Cienega Riparian Complex, Del Valle Riparian Complex, Piru Creek Riparian, Santa Clara River Riparian Shrubland	CNDDB	Indirect	Open canopy, deciduous riparian woodland close to water, along streams or wet meadows.). Reliant on groundwater-dependent riparian vegetation for breeding habitat (e.g., willows, alders, and cottonwoods). Typically eat insects. Occurrences along Santa Clara River and Piru Creek (CDFW 2019, eBird 2021, WFVZ 2020c).
Yellow-breasted chat Icteria virens	-/SSC	Likely	Cienega Riparian Complex, Del Valle Riparian Complex, Piru Creek Riparian	CNDDB, CAFSD	Indirect	Early-successional riparian habitats with a dense shrub layer and an open canopy. Occurrences along Santa Clara River (CNDDB 2019, eBird 2021, WFVZ 2020c). Occurrences along Santa Clara River and Piru Creek (CDFW 2019, eBird 2021).

<sup>&</sup>lt;sup>1</sup> Status codes:

Federal

= Listed as endangered under the federal Endangered Species Act

StateSE = Listed as Endangered under the California Endangered Species Act

FE = Listed as endangered under the federal Endangered Species Act
FT = Listed as threatened under the federal Endangered Species Act
SSC = CDFW species of special concern
SFP = CDFW fully protected species

FSS = Forest Service Sensitive Species

BLMS = Bureau of Land Management Sensitive Species

Likely: the species has documented occurrences and the habitat is high quality or quantity

Possible: no documented occurrences and the species' required habitat is moderate to high quality or quantity

Unlikely: no documented occurrences and the species' required habitat is of low to moderate quality or quantity

None: no potential to occur due to lack of habitat and/or the population is assumed extirpated

<sup>3</sup> Query source:

CAFSD: California Freshwater Species Database (TNC 2020)

CNDDB: California Natural Diversity Database (CDFW 2019)

eBird: (eBird 2021)

<sup>4</sup> Groundwater Dependent Ecosystem (GDE) association:

Direct: Species directly dependent on groundwater for some or all water needs

Indirect: Species dependent upon other species that rely on groundwater for some or all water needs

<sup>5</sup> Formerly Emys marmorata marmorata

Potential to Occur:

## Fish

Similar to the Fillmore Groundwater Basin, the Santa Clara River in the Piru Groundwater Basin likely supports limited fish populations because of the presence of sub-optimal water quality conditions (e.g., high temperature, low dissolved oxygen), poor habitat quality (e.g., low amounts of cover that provides refuge from predators and high flows), and the presence of non-native fish (Stoecker and Kelley 2005), although there is minimal literature regarding fish populations within these reaches. Piru Creek supports diverse native and non-native fish species assemblages because it has more instream shelter cover and more riparian vegetation than the Santa Clara River (Stoecker and Kelley 2005), which provide suitable spawning, feeding, and rearing habitat for fish. Disconnected ephemeral tributaries in the Piru Groundwater Basin can be used by fish species seasonally, but do not contain surface water year-round and are not connected to groundwater.

Four special-status fish species have the potential to occur in the interconnected reaches of the Piru Groundwater Basin (Table 4.1-4). An additional species, the Santa Ana Sucker, occurs in the Piru Groundwater Basin and is listed as threatened under the federal Endangered Species Act but those occurring in the Santa Clara River and tributaries have no special status due to uncertainties at the time of its listing regarding whether it is native to the Santa Clara River watershed. In the Piru Groundwater Basin, designated critical habitat for Southern California Steelhead includes the mainstem Santa Clara River upstream to the Piru Creek confluence and Piru Creek upstream to Santa Felicia Dam (Figure 4.1-8). *O. mykiss* use the mainstem Santa Clara River for passage to suitable spawning areas upstream to the Piru Creek confluence and Piru Creek. Most of the fish species listed in Table 4.1-4 are likely to occur in perennial reaches within the basin.

Table 4.1-4. Groundwater-dependent special-status fish with known occurrence or suitable habit in the Piru Groundwater Basin.

Common name Scientific name	Native or introduced	Status <sup>1</sup> Federal/ State	Occurrence in interconnected reaches	Source(s)	Habitat and occurrence within the Piru Groundwater Basin
Southern California steelhead Oncorhynchus mykiss irideus	Native	FE	Santa Clara River upstream to Piru Creek (likely migration, unlikely rearing), Piru Creek (likely migration, spawning, and rearing)	Howard et al. 2015, Howard and Booth 2016, United 2018, ACS 2002, Stoecker and Kelley 2005	Occur in freshwater systems and require adequate water conditions suitable for migration (i.e., flow, dissolved oxygen levels within the surface water, and water temperature suitable for passage) and suitable substrate (i.e., gravels) for spawning. Juvenile <i>O. mykiss</i> require suitable cover, flow, foraging conditions, and cool temperatures for rearing. Juvenile emigration (i.e., outmigration to the ocean) requires water conditions suitable for migration. <i>O. mykiss</i> migration (both upstream and downstream) can occur in all surface water reaches of the Piru Groundwater Basin when flows are sufficiently high (Stoecker and Kelley 2005). <i>O. mykiss</i> spawning and rearing likely occurs in Piru Creek. Rearing could occur in the Santa Clara River, but is unlikely due to poor habitat and temperature conditions (Stoecker and Kelley 2005).

Common name Scientific name	Native or introduced	Status <sup>1</sup> Federal/ State	Occurrence in interconnected reaches	Source(s)	Habitat and occurrence within the Piru Groundwater Basin
Pacific lamprey Entosphenus tridentatus	Native	SSC	Possible in the Santa Clara River in the Cienega and Del Valle Reaches (migration, pre-spawning holding, rearing).  Possible in Piru Creek (migration, pre-spawning holding, spawning, rearing)	UWCD 2018, Stoecker and Kelley 2005	Occur in freshwater systems and require adequate flows for migration, suitable substrate (i.e., gravels) for spawning, and adequate cover for pre-spawning holding. Juveniles (called ammocoetes) spend an extended period of time (between four and ten years) rearing while burrowed in sediments filter feeding on organic material and require suitable cover, flow, foraging conditions, and cool temperatures. Juvenile migrant (called macropthalmia) emigration (i.e., outmigration to the ocean) requires water conditions suitable for migration (i.e., water velocity and water depth, dissolved oxygen levels within the surface water, and water temperature suitable for passage). Pacific lamprey migration (both upstream and downstream) can occur in all surface water reaches of the Piru Groundwater Basin when flows are sufficiently high (Stoecker and Kelley 2005). Pacific lamprey spawning and rearing occurs in Piru Creek.
Unarmored threespine stickleback Gasterosteus aculeatus williamsoni	Native	SE, FE, SFP	Likely in the Santa Clara River and possible in Piru Creek	ACS 2002, Swift et al. 1993, CDFW 2019	Occurs in freshwater rivers and streams. Unarmored threespine stickleback populations have been documented within the interconnected mainstem Santa Clara River (CDFW 2019), likely occur within Piru Creek year-round, and likely use the disconnected tributary reaches when conditions are conducive to passage. Migration is largely localized and opportunistic; this species does not exhibit defined migration.

Common name Scientific name	Native or introduced	Status <sup>1</sup> Federal/ State	Occurrence in interconnected reaches	Source(s)	Habitat and occurrence within the Piru Groundwater Basin
Santa Ana sucker Catostomus santaanae	Native	FT (not in Santa Clara River Watershed) <sup>3</sup>	Likely in the Santa Clara River and Piru Creek	Howard and Booth 2016, United 2018, ACS 2002, Swift et al. 1993, CDFW 2019	Occurs in freshwater rivers and streams. The species has been observed within all surface water reaches (i.e., perennial tributaries, perennial mainstem, and ephemeral reaches) in the Piru Groundwater Basin (CDFW 2019, United 2018, Howard and Booth 2016). Recent genetics studies documented Santa Ana and Owens sucker hybrids ( <i>Catostomus santaanae</i> x <i>fumeiventris</i> ) within the Piru Groundwater Basin.
Arroyo chub Gila orcutti	Introduced (but native to other nearby watersheds)	SSC	Likely in Santa Clara River, Likely in Piru Creek	Howard et al. 2015, United 2018, ACS 2002, CDFW 2019	Occurs in freshwater rivers and streams. Although arroyo chub ( <i>Gila orcutti</i> ), a CDFW SSC, is not native to the Santa Clara River watershed, CDFW protects the species within the watershed. Arroyo chub has been observed in the perennial mainstem within the Piru Groundwater Basin and is likely to occur in the perennial tributary reaches and ephemeral reaches when conditions are conducive to passage (CDFW 2019, Howard et al. 2015, United 2018). Arroyo chub does not exhibit defined migration and the species' movement is largely localized and opportunistic.

Federal

FE = Listed as endangered under the federal Endangered Species Act SE = Listed as Endangered under the California Endangered Species Act

State

FT = Listed as threatened under the federal Endangered Species Act SSC = CDFW species of special concern FSS = Forest Service Sensitive Species SFP = CDFW fully protected species

BLMS = Bureau of Land Management Sensitive Species

Likely: the species has documented occurrences and the habitat is high quality or quantity

Possible: no documented occurrences and the species' required habitat is moderate to high quality or quantity

Unlikely: no documented occurrences and the species' required habitat is of low to moderate quality or quantity

*None:* no potential to occur due to lack of habitat and/or the population is assumed extirpated

<sup>&</sup>lt;sup>2</sup> Potential to Occur:

The Santa Ana sucker is federally threatened; however, because of previous uncertainty regarding whether it is native to the SCR watershed, United States Fish and Wildlife Service (USFWS) does not currently consider the species federally threatened within the SCR watershed (USFWS 2017).

## Fillmore

### Plants and natural communities

Four potentially groundwater dependent special-status plant species have been documented in the Fillmore Groundwater Basin (Table 4.1-5. One species, white rabbit-tobacco (*Pseudognaphalium leucocephalum*), was identified as likely to depend on groundwater. Multiple populations of white rabbit-tobacco have been observed in the open rock and sand wash bed of the Santa Clara River (East Grove Riparian Complex GDE Unit), and at least one population has been documented within more densely wooded areas. Two species were identified as possibly dependent on groundwater, umbrella larkspur (*Delphinium umbraculorum*) and Greata's aster (*Symphyotrichum greatae*), both of which were observed in the mountain range to the north of the basin and are associated with moist oak forests and cismontane and riparian woodland, respectively. The fourth species, late-flowered mariposa lily (*Calochortus fimbriatus*) is unlikely to depend on groundwater.

Ten potentially groundwater-dependent sensitive natural communities were documented with the potential to occur in the Fillmore Groundwater Basin (Table 4.1-5). Eight of these communities are identified as likely to depend on groundwater; these GDEs are predominantly within riparian habitats (e.g., black cottonwood woodland and forest, southern mixed riparian forest, southern riparian scrub) and occur in the East Grove Riparian Complex and Sespe Creek Riparian GDE units. Additional potentially groundwater-dependent sensitive natural communities include herbaceous communities associated with wetlands and marsh margins (e.g., hardstem and California bulrush marshes [Schoenoplectus (acutus, californicus) Herbaceous Alliance] and ashy ryegrass – creeping ryegrass turfs [Leymus cinereus – Leymus triticoides Herbaceous Alliance]) which were documented in the East Grove Riparian Complex.

Table 4.1-5. Special-status plant species and natural communities with known occurrence within the Fillmore Groundwater Basin.

Common name Scientific name	Status <sup>1</sup>	Association with GDE	Documented Occurrences in GDE units	Source <sup>2</sup>	Habitat and occurrence
Plants					
Late-flowered mariposa lily Calochortus fimbriatus	1B.3, S3, G3, not state or federally listed	Unlikely	Santa Clara River Riparian Shrubland, Tributary Riparian	CNDDB	Dry, open coastal and foothill woodland, chaparral; two CNDDB occurrences in the vicinity of Santa Paula Peak and Sespe Creek.
Umbrella larkspur Delphinium umbraculorum	1B.3, S3, G3, not state or federally listed	Possible	Tributary Riparian	CNDDB	Moist oak forest, foothill woodland; single CNDDB record from 1999 on the north side of Santa Paula Peak.
White rabbit-tobacco Pseudognaphalium leucocephalum	2B.2, S2, G4, not state or federally listed	Likely	Cienega Riparian Complex, East Grove Riparian Complex	CNDDB	Sandy or gravelly benches, dry stream or canyon bottoms; CNDDB records in sandy, rocky washes.
Greata's aster Symphyotrichum greatae	1B.3, S2, G2, not state or federally listed	Possible	Tributary Riparian	CNDDB	Mesic areas in broadleafed upland forest, chaparral, cismontane and riparian woodland; CNDDB records in Hopper and Pine canyons.
Sensitive Natural Communitie	S				
Ashy ryegrass – creeping ryegrass turfs  Leymus cinereus – Leymus triticoides herbaceous alliance	\$3, G3	Likely	East Grove Riparian Complex	VMSCR	Poorly drained floodplains, playas, drainage and valley bottoms, and marsh margins; occur throughout much of California at elevations below 10,000 ft.
Black cottonwood forest and woodland <i>Populus trichocarpa</i> forest and woodland alliance	S3, G5	Likely	Cienega Riparian Complex, East Grove Riparian Complex, Santa Clara River Riparian Shrubland, Sespe Creek Riparian	VMSCR	Seasonally flooded and permanently saturated soils on stream banks and alluvial terraces; occur throughout much of California except the Central Valley, Sacramento-San Joaquin Delta, and Mojave and Sonoran deserts.

Common name Scientific name	Status <sup>1</sup>	Association with GDE	Documented Occurrences in GDE units	Source <sup>2</sup>	Habitat and occurrence
California walnut groves  Juglans californica forest and woodland alliance	S3, G3	Unlikely	Sespe Creek Riparian, Tributary Riparian, Santa Clara River Riparian Shrubland	CalVeg, CNDDB, VMSCR	Riparian corridors, but most stands cover all hillslopes; occur along southern California coast and in Peninsular and Transverse ranges.
Fremont cottonwood forest and woodland Populus fremontii – Fraxinus velutina – Salix gooddingii forest and woodland alliance	S3, G4	Likely	Santa Clara River Riparian Shrubland	CalVeg, VMSCR	On floodplains, along low-gradient rivers and streams, and in alluvial fans and valleys with a dependable subsurface water supply; occur throughout much of California except the Sierra Nevada and Modoc Plateau.
Hardstem and California bulrush marshes Schoenoplectus (acutus, californicus) herbaceous alliance	S3S4, not globally ranked	Likely	East Grove Riparian Complex	VMSCR	Brackish to freshwater marshes, stream banks and bars of river mouth estuaries, ponds and lake shores, sloughs, and roadside ditches; occur throughout most of California at elevations below 8,200 ft.
Scale broom scrub  Lepidospartum squamatum  shrubland alliance	S3, G3	Unlikely	Santa Clara River Riparian Shrubland, Sespe Creek Riparian	CalVeg, VMSCR	Intermittently or rarely flooded, low-gradient alluvial deposits along streams, washes, and fans; occur in southern California in the inner central Coast Ranges, Transverse and Peninsular ranges, and the Mojave Desert.
Southern cottonwood willow riparian forest	S3.2, G3	Likely	Sespe Creek Riparian	CNDDB	Frequently flooded lands along rivers and streams; occur in the Transverse and Peninsular ranges from Santa Barbara County south and east to the edge of the deserts.
Southern mixed riparian forest	S2.1, G2	Likely	Tributary Riparian	CNDDB	Typically a younger successional stage of riparian forest due to disturbance or more frequent flooding.

Common name Scientific name	Status <sup>1</sup>	Association with GDE	Documented Occurrences in GDE units	Source <sup>2</sup>	Habitat and occurrence
Southern riparian scrub	S3.2, G3	Likely	East Grove Riparian Complex, Santa Clara River Riparian Shrubland	CNDDB	Sand and gravel bars close to river channels; occur along and at the mouths of perennial and intermittent streams of the southern Coast Ranges.
Southern willow scrub	S2.1, G3	Likely	Santa Clara River Riparian Shrubland	CNDDB	Generally on alluvium deposited near stream channels during floods; occur along major rivers of coastal southern California

<sup>1</sup> Status codes:

G = Global

T = Subspecies or variety

Federal

FT = Listed as threatened under the federal Endangered Species Act

FD = Federally delisted

State

S = Sensitive

SE = Listed as Endangered under the California Endangered Species Act

ST = Listed as Threatened under the California Endangered Species Act

SSC = CDFW species of special concern SFP = CDFW fully protected species

#### Rank

- 1 Critically Imperiled—At very high risk of extinction due to extreme rarity (often 5 or fewer populations), very steep declines, or other factors.
- 2 Imperiled—At high risk of extinction due to very restricted range, very few populations (often 20 or fewer), steep declines, or other factors.
- 3 Vulnerable At moderate risk of extinction or elimination due to a restricted range, relatively few populations (often 80 or fewer), recent and widespread declines, or other factors.
- 4 Apparently Secure Uncommon but not rare; some cause for long-term concern due to declines or other factors.
- 5 Demonstrably Secure Common; widespread and abundant.
- Q Taxonomic questions associated with this name

Ranks such as S2S3 indicate a ranking between S2 and S3

### California Rare Plant Rank (CRPR)

- 1B Plants rare, threatened, or endangered in California and elsewhere
- 2B Plants rare, threatened, or endangered in California, but more common elsewhere
- 4 More information needed about this plant, a review list
- 4 Plants of limited distribution, a watch list

#### CRPR Threat Ranks:

- 0.1 Seriously threatened in California (high degree/immediacy of threat)
- 0.2 Fairly threatened in California (moderate degree/immediacy of threat)
- 0.3 Not very threatened in California (low degree/immediacy of threats or no current threats known)
- <sup>2</sup> Sources: CNDDB (CDFW 2019), CalVeg (USDA 2014), VMSCR (Vegetation Map of the Santa Clara River; Stillwater Sciences 2019).

# Terrestrial and aquatic wildlife

Ten potentially groundwater-dependent special-status terrestrial and aquatic wildlife species were identified as having the potential to occur within the Fillmore Groundwater Basin: one amphibian species, seven bird species, and two reptile species —two reptile species and seven bird species (Table 4.1-6). Additional information on these species, including regulatory status, occurrences in GDE units, habitat associations, and documented occurrences in the groundwater basin is provided in Table 4.1-6.

Southwestern pond turtle (*Actinemys pallida*) and two-striped garter snake (*Thamnophis hammondii*) are the only wildlife species likely present (i.e., documented occurrences) in the Fillmore Groundwater Basin classified as directly groundwater dependent due to their association with riverine and lentic habitats. Southwestern pond turtles likely use habitat within the Santa Clara River GDE units (i.e., Santa Clara River Riparian Shrubland, Cienega Riparian Complex, and East Grove Riparian Complex) and Sespe Creek Riparian GDE for foraging (e.g., open water), nesting (e.g., grasses and forbs alliance), and/or overwintering (e.g., mixed riparian and willow scrub and cottonwood or willow alliance). Two-striped garter snake likely use freshwater (e.g., perennial riverine) and riparian (e.g., riparian mixed hardwood and/or shrub) habitats within the Santa Clara Riparian Shrubland, Sespe Creek Riparian GDEs for capturing prey. The Arroyo Toad was not documented within the Fillmore Groundwater Basin; however, suitable habitat within the East Grove Riparian Complex and Sespe Creek Riparian GDE units (e.g., freshwater wetland; riverwash; and cottonwood, willow, or mixed riparian alliances) could potentially support all the toad's life history stages (breeding, larval juvenile rearing, juvenile and adult foraging).

Indirectly groundwater-dependent bird species (i.e., Least Bell's vireo, southwestern willow flycatcher, western yellow-billed cuckoo [Coccyzus americanus occidentalis], white-tailed kite [Elanus leucurus], yellow warbler [Setophaga petechia], and yellow-breasted chat [Icteria virens]) use riparian habitat (e.g., willow/willow shrub, cottonwood, mixed riparian alliances) within the Santa Clara River GDE units (i.e., Santa Clara River Riparian Shrubland, Cienega Riparian Complex, and East Grove Riparian Complex) for foraging, nesting, and migrating. These GDE units include designated critical habitat for the southwestern willow flycatcher (Santa Clara River Riparian Shrubland [952 acres], Cienega Riparian Complex [116 acres], and East Grove Riparian Complex [923 acres]). The California condor's designated critical habitat is not located within a GDE unit; habitat use in the Fillmore Basin GDE Units (e.g., Sespe Creek Riparian) may be limited to foraging.

Table 4.1-6. Groundwater-dependent special-status terrestrial and aquatic wildlife species with known occurrence or suitable habit in the Fillmore Groundwater Basin.

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in Fillmore Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Fillmore Groundwater Basin
Amphibian	_			•		
Arroyo toad Anaxyrus californicus	FE/SSC	Possible	No documented occurrences	CAFSD	Direct	Washes, arroyos, sandy riverbanks, riparian areas with willows, sycamores, oaks, cottonwoods; needs exposed sandy streamsides with stable terraces for burrowing, with scattered vegetation for shelter, and areas of quiet water or pools free of predatory fishes with sandy or gravel bottoms without silt for breeding. The arroyo toad depends on groundwater for breeding in shallow still pools and riparian vegetation that provides foraging habitat (Rohde et al. 2019). It was historically found in the upper and lower Santa Clara River basin and currently persists in along Sespe Creek from Hot Springs Canyon upstream to the mouth of Tule Creek (Sweet 1992, as cited in USFWS 1999). Low potential to occur along the Santa Clara River (VCWPD and LADPW 2005).
Reptile						
Two-striped gartersnake Thamnophis hammondii	BLMS, FSS/SSC	Likely	Santa Clara Riparian Shrubland, Sespe Creek Riparian	CNDDB, CAFSD	Direct	Highly aquatic snake species. Found in or near permanent fresh water, often along streams with rocky beds and riparian vegetation. Prey includes fish, fish eggs, tadpoles, newt larvae, small frogs and toads, leeches, and earthworms. Commonly found in the Santa Clara River Watershed (UWCD 2018). Occurrences along Santa Clara River bank on the south edge of Fillmore and on Sespe Creek (<1 mile upstream of the Sespe Creek Riparian GDE) unit in 2011 (CDFW 2019).

Common name Scientific name	Status¹ Federal/State	Potential to occur in Fillmore Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Fillmore Groundwater Basin
Southwestern pond turtle <i>Actinemys</i> pallida <sup>5</sup>	BLMS, FSS/SSC	Likely	East Grove Riparian Complex, Sespe Creek Riparian	CNDDB, CAFSD	Direct	Ponds, lakes, rivers, streams, creeks, marshes, and irrigation ditches with basking sites. Feeds on aquatic plants, invertebrates, worms, frog and salamander eggs and larvae, crayfish, and occasionally frogs and fish. Relies on surface water that may be supported by groundwater (Rhode et al. 2019). Documented throughout the Santa Clara Rivera and Sespe Creek watersheds. Occurrences include Santa Clara River approximately 3 miles downstream of the confluence with Sespe Creek in 2016 and Sespe Creek between 2 and 4 miles north of Fillmore (CDFW 2019).
Bird						
California condor Gymnogyps californianus	FE/SE	Likely	Sespe Creek Riparian	CNDDB	Indirect	Require vast expanses of open savannah, grasslands, and foothill chaparral in mountain ranges of moderate altitude; deep canyons containing clefts in rocky walls and large trees provide nest sites; forages up to 100 miles from roost to nest. Forage in grasslands, oak savanna habitats, and Condors may rely on groundwater-dependent vegetation for nesting in foothill grasslands, oak savanna habitats, and old-growth forest (Rohde et al. 2019). Documented near Sespe Creek at Grand Avenue terminus (eBird 2021). One acre of USFWS critical habitat located within the Fillmore Groundwater Basin (USFWS 1976).

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in Fillmore Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Fillmore Groundwater Basin
Least Bell's vireo Vireo bellii pusillus	FE/SE	Likely	Cienega Riparian Complex, East Grove Riparian Complex, Santa Clara River Riparian Shrubland	CNDDB, CAFSD	Indirect	Nests in dense vegetative cover of riparian areas; often nests in willow or mulefat; forages in dense, stratified canopy. This species relies on groundwater-dependent vegetation in riparian areas, particularly during breeding periods (Rohde et al. 2019). Eats insects, fruits, and berries. Documented throughout the Santa Clara River and Sespe Creek in the Fillmore Groundwater Basin (CDFW 2019, eBird 2021, WFVZ 2020c). Documented breeding occurrences in: the TNC properties listed, Sespe Cienega (Cienega Riparian Complex) (eBird 2021), the Taylor property and the Hedrick Ranch Nature Area (East Grove Riparian Complex); CDFW's Cienaga Springs Ecological Preserve (Cienega Riparian Complex and Santa Clara River Riparian Shrubland) (WFVZ 2020a, WFVZ 2020b).
Southwestern willow flycatcher Empidonax traillii extimus	FE/SE	Likely	Cienega Riparian Complex, East Grove Riparian Complex, Santa Clara River Riparian Shrubland	CNDDB, CAFSD	Indirect	Dense brushy thickets within riparian woodland often dominated by willows and/or alder, near permanent standing water. Reliant on groundwater-dependent riparian vegetation, including for nest sites that are typically located near slow-moving streams, or side channels and marshes with standing water and/or wet soils (Rohde et al. 2019). Feeds on insects, fruits, and berries. Occurrences throughout the Santa Clara River (CDFW 2019, eBird 2021). Critical habitat located along the Santa Clara River in the East Grove Riparian Complex, Santa Clara River, and Cienega Riparian Complex GDE units (USFWS 2013).

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in Fillmore Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Fillmore Groundwater Basin
Western yellow- billed cuckoo Coccyzus americanus occidentalis	BLMS, FSS, FT/SE	Likely	East Grove Riparian Complex, Santa Clara River Riparian Shrubland	CNDDB	Indirect	Summer resident of valley foothill and desert riparian habitats; nests in open woodland with clearings and low, dense, scrubby vegetation. Reliant on groundwater-dependent riparian vegetation for habitat (Rhode et al. 2019). Occurrences along Santa Clara River in the East Grove Riparian Complex GDE (CDFW 2019, eBird 2021), and in TNC's Hedrick Ranch Nature Area (East Grove Riparian Complex) (WFVZ 2020b). Historical populations documented along Sespe Creek west of Fillmore in 1924 are presumed extant (CDFW 2019).
White-tailed kite Elanus leucurus	BLMS/SFP	Likely	Cienega Riparian Complex, East Grove Riparian Complex, Santa Clara River Riparian Shrubland	CNDDB	Indirect	Lowland grasslands and wetlands with open areas; nests in trees near open foraging area. Predominately preys on small mammals, but its diet also includes birds, and lizards. Occurrences along Santa Clara River (CDFW 2019, eBird 2021, WFVZ 2020c). Breeding documented in CDFW's Cienaga Springs Ecological Preserve (Cienega Riparian Complex and Santa Clara River Riparian Shrubland) (WFVZ 2020a).
Yellow warbler Setophaga petechia	-/SSC	Likely	Cienega Riparian Complex, East Grove Riparian Complex, Santa Clara River Riparian Shrubland, Sespe Creek Riparian,	CNDDB	Indirect	Open canopy, deciduous riparian woodland close to water, along streams or wet meadows.). Reliant on groundwater-dependent riparian vegetation for breeding habitat (e.g., willows, alders, and cottonwoods). Typically eat insects. Occurrences along Santa Clara River (CNDDB 2019, eBird 2021, WFVZ 2020c) and near Sespe Creek at Grand Avenue terminus (CNDDB 2019, eBird 2021). Breeding documented in CDFW's Cienaga Springs Ecological Preserve (Cienega Riparian Complex) (WFVZ 2020a), and TNC's Taylor property and the Hedrick Ranch Nature Area (East Grove Riparian Complex) (WFVZ 2020b).

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in Fillmore Groundwater Basin <sup>2</sup>	Documented occurrences in GDE units	Query source <sup>3</sup>	GDE association <sup>4</sup>	Habitat and documented occurrences in Fillmore Groundwater Basin
Yellow-breasted chat Icteria virens	-/SSC	Likely	Cienega Riparian Complex, East Grove Riparian Complex, Santa Clara River Riparian Shrubland	CNDDB, CAFSD	Indirect	Early-successional riparian habitats with a dense shrub layer and an open canopy. Occurrences along Santa Clara River (CDFW 2019, eBird 2021, WFVZ 2020c). Breeding documented in CDFW's Cienaga Springs Ecological Preserve (Cienega Riparian Complex) (WFVZ 2020a), and suspected breeding in TNC's Hedrick Ranch Nature Area (East Grove Riparian Complex) (WFVZ 2020b).

<sup>&</sup>lt;sup>1</sup> Status codes:

Federal State

FE = Listed as endangered under the federal Endangered Species Act SE = Listed as Endangered under the California Endangered Species Act

FT = Listed as threatened under the federal Endangered Species Act SSC = CDFW species of special concern SSS = Forest Service Sensitive Species SFP = CDFW fully protected species

Likely: the species has documented occurrences and the habitat is high quality or quantity

Possible: no documented occurrences and the species' required habitat is moderate to high quality or quantity

Unlikely: no documented occurrences and the species' required habitat is of low to moderate quality or quantity

None: no potential to occur due to lack of habitat and/or the population is assumed extirpated

3 Ouerv source

CAFSD: California Freshwater Species Database (TNC 2020)

CNDDB: California Natural Diversity Database (CDFW 2019)

eBird: (eBird 2021)

Direct: Species directly dependent on groundwater for some or all water needs

Indirect: Species dependent upon other species that rely on groundwater for some or all water needs

<sup>5</sup> Formerly *Emys marmorata marmorata* 

BLMS = Bureau of Land Management Sensitive Species

<sup>&</sup>lt;sup>2</sup> Potential to Occur:

<sup>&</sup>lt;sup>4</sup> Groundwater Dependent Ecosystem (GDE) association:

## Fish

Fish are dependent on groundwater if they use interconnected surface water as part of their life cycle including spawning, rearing, and migration. As discussed in Section 3.3 interconnected surface waters in the Fillmore Groundwater Basin occur in along the East Grove Riparian Complex and Cienega Riparian Complex GDE units of the Santa Clara River and likely in portions of Sespe Creek.

The mainstem Santa Clara River likely support limited fish populations because of the presence of sub-optimal water conditions (e.g., high temperature, low dissolved oxygen), poor habitat quality (e.g., low amounts of cover that provides refuge from predators and high flows), and the presence of non-native fish (Stoecker and Kelley 2005), although there is minimal literature regarding fish populations within these reaches. Sespe Creek supports diverse native and non-native fish species assemblages it has more instream shelter cover and more riparian vegetation than the Santa Clara River (Stoecker and Kelley 2005), which provide suitable spawning, feeding, and rearing habitat for fish. Disconnected ephemeral tributaries in the Fillmore Groundwater Basin can be used by fish species seasonally, but do not contain surface water year-round and are not connected to groundwater and thus not considered here.

Four special-status fish species have the potential to occur in the interconnected reaches of the Fillmore Groundwater Basin (Table 4.1-7). An additional species, the Santa Ana Sucker, occurs in the Fillmore Groundwater Basin and is listed as threatened under the federal Endangered Species Act but those occurring in the Santa Clara River and tributaries have no special status due to uncertainties at the time of its listing regarding whether it is native to the Santa Clara River watershed. In the Fillmore Groundwater Basin, the National Marine Fisheries Service (NMFS) designated critical habitat for Southern California Steelhead in the mainstem Santa Clara River and Sespe Creek (Figure 4.1-6). Due to the life history complexities, steelhead is the term used to describe the anadromous life history type and *O. mykiss* is generally used when life history types are indistinguishable, such as when they are juveniles, or when referring in general terms to a population whose life history type expression is uncertain. *O. mykiss* utilize the mainstem Santa Clara River for passage to suitable spawning areas in Sespe Creek and further upstream to Piru Creek. Resident fish likely use perennial reaches year-round and intermittent reaches when water is present.

Table 4.1-7. Groundwater-dependent special-status fish with known occurrence or suitable habit in the Fillmore Groundwater Basin.

Common name Scientific name	Native or introduced	Status <sup>1</sup> Federal/ State	Occurrence in interconnected reaches	Source(s)	Habitat and occurrence within the Fillmore Groundwater Basin
Southern California steelhead Oncorhynchus mykiss irideus	Native	FE	Santa Clara River (migration likely, rearing unlikely), Sespe Creek (likely migration, spawning, and rearing)	Howard et al. 2015, Howard and Booth 2016, United 2018, ACS 2002, Stoecker and Kelley 20005	Occur in freshwater systems and require adequate water conditions suitable for migration (i.e., flow, dissolved oxygen levels within the surface water, and water temperature suitable for passage) and suitable substrate (i.e., gravels) for spawning. Juvenile <i>O. mykiss</i> require suitable cover, flow, foraging conditions, and cool temperatures for rearing. Juvenile emigration (i.e., outmigration to the ocean) requires water conditions suitable for migration. <i>O. mykiss</i> migration (both upstream and downstream) can occur in all surface water reaches of the Fillmore Groundwater Basin when flows are sufficiently high (Stoecker and Kelley 2005). <i>O. mykiss</i> spawning and rearing occurs in Sespe Creek. Rearing is unlikely in the Santa Clara River due to poor habitat and temperature conditions (Stoecker and Kelley 2005).

Common name Scientific name	Native or introduced	Status <sup>1</sup> Federal/ State	Occurrence in interconnected reaches	Source(s)	Habitat and occurrence within the Fillmore Groundwater Basin
Pacific lamprey Entosphenus tridentatus	Native	SSC	Santa Clara River (likely migration, pre-spawning holding, rearing), Sespe Creek (likely migration, pre-spawning holding, spawning, rearing)	United 2018, Stoecker and Kelley 2005	Occur in freshwater systems and require adequate flows for migration, suitable substrate (i.e., gravels) for spawning, and adequate cover for pre-spawning holding. Juveniles (called ammocoetes) spend an extended period of time (between four and ten years) rearing while burrowed in sediments filter feeding on organic material and require suitable cover, flow, foraging conditions, and cool temperatures. Juvenile migrant (called macropthalmia) emigration (i.e., outmigration to the ocean) requires water conditions suitable for migration (i.e., water velocity and water depth, dissolved oxygen levels within the surface water, and water temperature suitable for passage). Pacific lamprey migration (both upstream and downstream) is likely in all surface water reaches of the Fillmore Groundwater Basin when flows are sufficiently high (Stoecker and Kelley 2005). Pacific lamprey spawning and rearing occurs in Sespe Creek.
Unarmored threespine stickleback Gasterosteus aculeatus williamsoni	Native	SE, FE, SFP	Likely in the Santa Clara River, possible in Sespe Creek	ACS 2002, Swift et al. 1993, CDFW 2019	Occurs in freshwater rivers and streams. Unarmored threespine stickleback populations have been documented within the interconnected mainstem Santa Clara River (CDFW 2019), likely occur within Sespe Creek year-round, and likely use the disconnected tributary reaches when conditions are conducive to passage. Migration is largely localized and opportunistic; this species does not exhibit defined migration.

Common name Scientific name	Native or introduced	Status <sup>1</sup> Federal/ State	Occurrence in interconnected reaches	Source(s)	Habitat and occurrence within the Fillmore Groundwater Basin
Santa Ana sucker Catostomus santaanae	Native	FT (not in Santa Clara River Watershed) <sup>3</sup>	Likely in the Santa Clara River and Sespe Creek	Howard and Booth 2016, United 2018, ACS 2002, Swift et al. 1993, CDFW 2019	Occurs in freshwater rivers and streams. The species occurs within all surface water reaches in the Fillmore Groundwater Basin (CDFW 2019, United 2018, Howard and Booth 2016). Recent genetics studies reveal the presence of the Santa Ana and Owens sucker hybrid ( <i>Catostomus santaanae x fumeiventris</i> ) within Fillmore Groundwater Basin.
Arroyo chub Gila orcutti	Introduced (but native to other nearby watersheds)	SSC	Likely in the Santa Clara River, possible in Sespe Creek	Howard et al. 2015, United 2018, ACS 2002, CDFW 2019	Occurs in freshwater rivers and streams. Although Arroyo chub ( <i>Gila orcutti</i> ), a CDFW SSC, is not native to the Santa Clara River watershed, CDFW protects the species within the watershed. Arroyo chub occurs in the perennial mainstem within Fillmore and is likely to occur in the perennial tributary reaches and ephemeral reaches when conditions are conducive to passage (CDFW 2019, Howard et al. 2015, United 2018). Arroyo chub does not exhibit defined migration and the species' movement is largely localized and opportunistic.

1 Status codes: Federal

FΕ = Listed as endangered under the federal Endangered Species Act

= Listed as threatened under the federal Endangered Species Act

= Forest Service Sensitive Species

BLMS = Bureau of Land Management Sensitive Species

State

= Listed as Endangered under the California Endangered Species Act SE

= CDFW species of special concern SFP

= CDFW fully protected species

Likely: the species has documented occurrences and the habitat is high quality or quantity

Possible: no documented occurrences and the species' required habitat is moderate to high quality or quantity

Unlikely: no documented occurrences and the species' required habitat is of low to moderate quality or quantity

None: no potential to occur due to lack of habitat and/or the population is assumed extirpated

Potential to Occur:

<sup>3</sup> The Santa Ana sucker is federally threatened; however, because of previous uncertainty regarding whether it is native to the SCR watershed, United States Fish and Wildlife Service (USFWS) does not currently consider the species federally threatened within the SCR watershed (USFWS 2017)

# 4.2 Invasive Species

Non-native and invasive species are distributed throughout the Santa Clara River watershed, including the Fillmore and Piru groundwater basins. Invasive species have a negative impact on the riparian corridor and threaten native species populations. Two invasive plant species, Arundo (giant reed; *Arundo donax*) and saltcedar (*Tamarix* spp.), are widely distributed within the Fillmore and Piru groundwater basins. Arundo and salt cedar were included as vegetation community types (see Section 4.1.1). The extent of mapped Arundo and salt cedar are shown in Figure 4.2.-1 and acreages of community are presented in Appendix A.

Arundo is a highly aggressive, naturalized landscape plant that invades riparian zones by establishing dense, monospecific clonal stands (DiTomaso and Healy 2007). It spreads quickly and severely impacts the ecology of the riparian corridor (Stillwater Sciences and URS 2007) and uses a large amount of water to fuel its growth (Bell 1997, Geissow et al. 2011). In California, giant reed is known to increase the risk of flooding, create fire hazards, outcompete indigenous plant species for scarce water resources, and reduce the value of riparian habitat for wildlife (Bell 1994, Bell 1997, DiTomaso 1998). The least Bell's vireo and other riparian birds require structural diversity provided by riparian scrub and mature forest communities for breeding (Zembal 1990, Bell 1994, Bell 1997). When natural riparian vegetation types are replaced by thick stands of giant reed, bird species abundance and other native wildlife have been found to decline (Bell 1994, Bell 1997, Herrera and Dudley 2003, Kisner 2004, Labinger and Greaves 2001).

In general, invading saltcedar lowers wildlife habitat value in riparian ecosystems heavily by decreasing available food sources and altering structural characteristics (Shafroth et al. 2005). Monotypic stands provide limited cover for large mammals, and fewer nesting sites for birds and herpetofauna in more southern latitudes due to lack of shading in mid- to late-summer (Hunter et al. 1988, Lovich and DeGouvenain 1998, Shafroth et al. 2005). Both the endangered southwestern willow flycatcher and the candidate for federal endangered species list western yellow-billed cuckoo prefer native forests in some cases, but incorporate some habitat with saltcedar into their breeding territory (Shafroth et al. 2005).

Invasive amphibian species, American bullfrog (*Lithobates catesbeianus*) and African clawed frog (*Xenopus laevis*), are documented on the Santa Clara River, including near the confluence with lower Sespe Creek and Piru Creek (Santa Clara River Trustee Council 2008). These amphibian species are found within or adjacent to aquatic habitat, including ponds, streams, reservoirs, and lakes. Both species likely prey on native species and have negative impacts on native amphibian species (e.g., Arroyo toad).

Many non-native fish species including black bullhead (*Ameiurus melas*), various sunfish species (*Lepomis* sp.) (e.g., green sunfish [*Lepomis cyanellus*] and bluegill [*Lepomis macrochirus*]), and bass species (*Micropterus* sp.) (e.g., largemouth bass [*Micropterus salmoides*]), have been documented within the Piru and Fillmore perennial tributaries year-round or within the other reaches within the SCR watershed (Stoecker and Kelley 2005). The distribution of invasive species within the mainstem and tributaries is not fully known; however, these species are either likely to occur or have the potential to occur within the perennial mainstem and ephemeral reaches due to the species' observation throughout the SCR watershed. Non-native predatory fish may have a large impact on native fish populations (e.g., salmonids), reducing the size of already diminished populations and limiting their ability to recover in response to habitat restoration

efforts. In the NMFS (2012) Southern California steelhead recovery plan non-native species were designated a "very high threat" to the steelhead population in the Santa Clara River.




Figure 4.2-1. Invasive species mapped within the Fillmore and Piru groundwater basins.

## 4.3 Ecological Value

The ecological value of each GDE unit was characterized by evaluating the presence and groundwater-dependence of special-status species and ecological communities, and the vulnerability of these species and their habitat to changes in groundwater levels (Rohde et al. 2018). In addition, the presence of natural or near-natural conditions and ecosystem function was also considered.

#### 4.3.1 Piru

## Del Valle Riparian Complex GDE Unit

The Del Valle Riparian Complex GDE Unit was determined to have high ecological value because: (1) it supports a relatively large number of special-status species and ecological communities (Tables 4.1-2, 4.1-3, and 4.1-4), (2) contains 433 acres of designated critical habitat for Southwestern willow flycatcher and 436 acres of designated critical habitat for least Bell's vireo (Figure 4.1-8), (3) supports native special-status species and natural communities with a known or high likelihood of direct groundwater dependence (two plants, three natural communities, two reptiles, and one fish; Tables 4.1-2, 4.1-3, and 4.1-4), and (4) includes species and ecological communities that are highly or moderately vulnerable to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). The reach of the Santa Clara River in the Del Valle Riparian Complex GDE Unit is considered perennial and is typically connected to groundwater. The degree to which interconnected surface waters in this reach are maintained by releases from upstream effluent sources is unknown, but is believed to be significant. It provides important habitat for special-status aquatic and semi-aquatic species potentially including Southwestern pond turtle, two-striped garter snake, unarmored threespine stickleback, and arroyo chub and this habitat is vulnerable to groundwater uses that reduce the amount and quality of riverine habitat. Southern California steelhead and Pacific lamprey are not known to occur in or upstream of this reach. The Del Valle Riparian Complex GDE Unit contains 34% of the total GDE acreage in the Piru Groundwater Basin (Table 2.3-2).

#### Santa Clara River Riparian Shrubland GDE Unit

The Santa Clara River Riparian Shrubland GDE Unit was determined to have **moderate ecological value** because: (1) it supports a moderate number of special-status species and ecological communities (Tables 4.1-2, 4.1-3, and 4.1-4), (2) contains 317 acres of designated critical habitat for Southwestern willow flycatcher and approximately 3.8 miles of designated critical habitat for Southern California steelhead (Figure 4.1-8), (3) supports native special-status species and natural communities with a known or high likelihood of direct groundwater dependence (three natural communities and three fish; Tables 4.1-2, 4.1-3, and 4.1-4), and (4) includes species and ecological communities that are somewhat vulnerable to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). The Santa Clara River in this GDE unit is considered intermittent and is not connected to groundwater. It may provide migration habitat for anadromous species (i.e., Southern California steelhead, Pacific lamprey) but this habitat has low vulnerability to groundwater reduction because most migration occurs during seasonal high flow periods. The unit contains 25% of the total GDE acreage in the Piru Groundwater Basin (Table 2.3-2).

#### Cienega Riparian Complex GDE Unit

The Cienega Riparian Complex GDE Unit was determined to have high ecological value because: (1) it supports a moderate number of special-status species and ecological communities (Tables 4.1-2, 4.1-3, and 4.1-4), (2) contains 154 acres of designated critical habitat for Southwestern willow flycatcher and approximately 0.2 miles of designated critical habitat for Southern California steelhead (Figure 4.1-8), (3) supports native special-status species and natural communities with a known or high likelihood of direct groundwater dependence (one plant, two reptiles, and four fish; Tables 4.1-2, 4.1-3, and 4.1-4), and (4) includes species and ecological communities that are highly or moderately vulnerable to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). The reach of the Santa Clara River in the Cienega Riparian Complex GDE Unit is considered perennial and is connected to groundwater except during droughts. It provides important habitat for special-status aquatic and semi-aquatic species potentially including Southwestern pond turtle, two-striped garter snake, Southern California steelhead, Pacific lamprey, unarmored threespine stickleback, and arroyo chub, and this habitat is vulnerable to groundwater uses that reduce the amount and quality of riverine habitat. The unit contains 12% of the total GDE acreage in the Piru Groundwater Basin (Table 2.3-2).

#### Piru Creek Riparian GDE Unit

The Piru Creek Riparian GDE Unit was determined to have **high ecological value** because: (1) it supports a relatively high number of special-status species and ecological communities (Tables 4.1-2, 4.1-3, and 4.1-4), (2) contains 246 acres of designated critical habitat for Southwestern willow flycatcher and approximately 4.9 miles of designated critical habitat for Southern California steelhead (Figure 4.1-8), (3) supports native special-status species and natural communities with a known or high likelihood of direct groundwater dependence (one plant, five natural communities, two reptiles, and five fish; Tables 4.1-2, 4.1-3, and 4.1-4), and (4) includes species and ecological communities that are highly or moderately vulnerable to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). Piru Creek in this GDE unit is considered perennial, though baseflows are maintained by releases from Santa Felicia Dam which likely also raises the groundwater level in this area. The lower portion of Piru Creek near the confluence with the Santa Clara River periodically lacks surface flow. Piru Creek provides important habitat for special-status aquatic and semi-aquatic species potentially including Southwestern pond turtle, two-striped garter snake, Southern California steelhead, Pacific lamprey, unarmored threespine stickleback, and arroyo chub. Because Piru Creek's is not generally connected with groundwater, this habitat is not vulnerable to groundwater uses that reduce the amount and quality of stream habitat. The unit contains 25% of the total GDE acreage in the Piru Groundwater Basin (Table 2.3-2).

## Tributary Riparian GDE Unit

The Tributary Riparian GDE Unit was determined to have **moderate ecological value** because: (1) it supports a relatively low number of special-status species and ecological communities (Tables 4.1-2 and 4.1-3), (2) contains 5.6 acres of designated critical habitat for least Bell's vireo (Figure 4.1-8), (3) supports few native special-status species and natural communities with a likely or possible groundwater dependence (three plants and one natural community; Tables 4.1-2 and 4.1-3), and (4) primarily includes species and ecological communities with little vulnerability to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). They do, however, support valuable

riparian habitat and likely movement corridors for a variety of native wildlife species. The tributary streams in this GDE unit are considered ephemeral and are not interconnected with groundwater. Typically, these streams only support surface flow immediately after storm events and thus provide little habitat value for fish and other aquatic species. The unit contains 5% of the total GDE acreage in the Piru Groundwater Basin (Table 2.3-2).

#### 4.3.2 Fillmore

#### Santa Clara River Riparian Shrubland GDE Unit

The Santa Clara River Riparian Shrubland GDE Unit was determined to have moderate ecological value because: (1) it supports a relatively high number of special-status species and ecological communities (Tables 4.1-2, 4.1-3, and 4.1-4), (2) contains 952 acres of designated critical habitat for Southwestern willow flycatcher and approximately 7.6 miles of designated critical habitat for Southern California steelhead (Figure 4.1-3), (3) supports native special-status species and natural communities with a known or high likelihood of direct groundwater dependence (four natural communities, one reptile, and two fish; Tables 4.1-2, 4.1-3, and 4.1-4), and (4) includes species and ecological communities that are somewhat vulnerable to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). The Santa Clara River in this GDE unit is considered intermittent and is not connected to groundwater or to upstream and downstream perennial reaches in most years (Figure 3.1-1). It provides migration habitat for special-status anadromous species (i.e., Southern California steelhead, Pacific lamprey) but this habitat has low vulnerability to groundwater reduction because most migration occurs during seasonal high flow periods. The unit contains a relatively large amount of GDE area (1,046 acres), accounting for 40% of the total GDE acreage in the Fillmore Groundwater Basin (Table 2.3-2).

## Cienega Riparian Complex GDE Unit

The Cienega Riparian Complex GDE Unit was determined to have high ecological value because: (1) it supports a relatively large number of special-status species and ecological communities (Tables 4.1-5, 4.1-6, and 4.1-7), (2) contains 116 acres of designated critical habitat for Southwestern willow flycatcher and approximately 1.1 miles of designated critical habitat for Southern California steelhead (Figure 4.1-8), (3) supports native special-status species and natural communities with a known or high likelihood of direct groundwater dependence (one plant, one natural community, and four fish; Tables 4.1-5, 4.1-6, and 4.1-7), and (4) includes species and ecological communities that are highly or moderately vulnerable to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). The reach of the Santa Clara River in the Cienega Riparian Complex GDE Unit is considered perennial and is connected to groundwater except during droughts. It provides important habitat for special-status aquatic and semi-aquatic species potentially including Southwestern pond turtle, Southern California steelhead, Pacific lamprey, unarmored threespine stickleback, and arroyo chub, and this habitat is vulnerable to groundwater uses that reduce the amount and quality of riverine habitat. The unit contains 5% of the total GDE acreage in the Fillmore Groundwater Basin (Table 2.3-2).

## East Grove Riparian Complex GDE Unit

The East Grove Riparian Complex GDE Unit was determined to have **high ecological value** because: (1) it supports a relatively large number of special-status species and ecological communities (Tables 4.1-5, 4.1-6, and 4.1-7), (2) contains 923 acres of designated critical habitat

for Southwestern willow flycatcher and approximately 3.2 miles of designated critical habitat for Southern California steelhead (Figure 4.1-8), (3) supports native special-status species and natural communities with a known or high likelihood of direct groundwater dependence (one plant, four natural communities, one reptile, and three fish; Tables 4.1-5, 4.1-6, and 4-1-7), and (4) includes species and ecological communities that are highly or moderately vulnerable to changes in groundwater discharge or groundwater levels that could substantially alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). The reach of the Santa Clara River in the East Grove Riparian Complex GDE Unit is considered perennial and is typically connected to groundwater. It provides important habitat for special-status aquatic and semi-aquatic species potentially including Southwestern pond turtle, Southern California steelhead, Pacific lamprey, unarmored threespine stickleback, and arroyo chub and this habitat is vulnerable to groundwater uses that reduce the amount and quality of riverine habitat. The unit contains a relatively large amount of GDE area (1,101.9 acres), accounting for 43% of the total GDE acreage in the Fillmore Groundwater Basin (Table 2.3-2).

## Tributary Riparian GDE Unit

The Tributary Riparian GDE Unit was determined to have **moderate ecological value** because: (1) it supports a relatively low number of special-status species and ecological communities (Tables 4.1-5, 4.1-6, and 4.1-7), (2) contains no designated critical habitat (Figure 4.1-8), (3) supports few native special-status species and natural communities with a likely or possible dependence on groundwater (two plants and one natural community; Table 4.1-5), and (4) primarily includes species and ecological communities with little vulnerability to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). They do, however, support valuable riparian habitat and likely movement corridors for a variety of native wildlife species. The tributary streams in this GDE unit are considered ephemeral and are not connected to groundwater. Typically, these streams only support surface flow immediately after storm events and thus provide little habitat value for fish and other aquatic species. The Tributary Riparian GDE Unit contains 8% of the total GDE acreage in the Fillmore Groundwater Basin (Table 2.3-2).

#### Sespe Creek Riparian Complex GDE Unit

The Sespe Creek Riparian Complex GDE Unit was determined to have moderately high ecological value because: (1) it supports a moderate number of special-status species and ecological communities (Tables 4.1-5, 4.1-6, and 4.1-7), (2) contains approximately 3.2 miles of designated critical habitat for Southern California steelhead (Figure 4.1-8), (3) supports native special-status species and natural communities with a known or high likelihood of direct groundwater dependence (two natural communities, two reptiles, and three fish; Tables 4.1-5, 4.1-6, and 4.1-7), and (4) includes species and ecological communities that are highly or moderately vulnerable to changes in groundwater discharge or groundwater levels that could alter their distribution, species composition, and/or health (Rohde et al. 2018, 2019). The upper 2 miles of Sespe Creek in this GDE unit are considered perennial, while the lower portion of Sespe Creek is likely intermittent (Figure 3.1-1). Sespe Creek's connection to groundwater is undetermined. Sespe Creek provides important habitat for special-status aquatic and semi-aquatic species likely including Southwestern pond turtle, two-striped garter snake, Southern California steelhead, Pacific lamprey, unarmored threespine stickleback, and arroyo chub, and this habitat is vulnerable to groundwater uses that reduce the amount and quality of stream habitat. The unit contains 4% of the total GDE acreage in the Fillmore Groundwater Basin (Table 2.3-2).

# 5 POTENTIAL EFFECTS OF GROUNDWATER MANAGEMENT ON GDFS

This section presents the methods and results of our analysis to identify how groundwater management could affect GDEs in the Fillmore and Piru groundwater basins. Adverse effects (impacts) on GDEs are considered undesirable results under SGMA (State of California 2014). The analysis is based on the hydrologic conditions affecting GDEs and their susceptibility to changing groundwater conditions, trends in biological condition of the GDEs, and climate change projections and other anticipated conditions or management actions likely to affect GDEs in the future.

## 5.1 Approach

SGMA describes six groundwater conditions that could cause undesirable results, including adverse impacts on GDEs. These are (1) chronic lowering of groundwater levels, (2) reduction of groundwater storage, (3) seawater intrusion, (4) degraded groundwater quality, (5) land subsidence, and (6) depletion of interconnected surface waters. Rohde et al. (2018) identify chronic lowering of groundwater levels, degraded water quality, and depletions of interconnected surface water as the most likely conditions to have direct effects on GDEs, potentially leading to an undesirable result. Following this guidance and based on available information for the Fillmore and Piru groundwater basins, we have eliminated reduction of groundwater storage, seawater intrusion (the subbasins are not located in close proximity to the ocean), and land subsidence from consideration because they are not relevant to GDEs in the Fillmore and Piru groundwater basins.

We evaluated the potential for chronic lowering of groundwater levels, degraded groundwater quality, and depletion of interconnected surface waters to cause direct effects on GDEs compared to baseline conditions, with a focus on effects related to groundwater levels. First, we identified baseline hydrologic conditions for the GDE units using available information (Section 1.2 and Section 3). Next, we determined each GDE unit's susceptibility to changing groundwater conditions using available hydrologic data, climate change projections, and the GDE susceptibility classifications (Rohde et al. 2018) summarized in Table 5.1-1.

**Table 5.1-1.** Susceptibility classifications developed for evaluation of a GDE's susceptibility to changing groundwater conditions (Rohde et al. 2018).

Susceptibility classifications	
High Susceptibility	Current groundwater conditions for the selected hydrologic data fall outside the baseline range. <sup>1</sup>
Moderate Susceptibility	Current groundwater conditions for the selected hydrologic data fall within the baseline range but future changes in groundwater conditions are likely to cause it to fall outside the baseline range. The future conditions could be due to planned or anticipated activities that increase or shift groundwater production, causing a potential effect on a GDE.
Low Susceptibility	Current groundwater conditions for the selected hydrologic data fall within the baseline range and no future changes in groundwater conditions are likely to cause the hydrologic data to fall outside the baseline range.

<sup>1</sup> For purposes of this analysis the baseline range is defined as the range of variability of the shallow groundwater depth for the period of record through 2015, with a minimum of 10 years (2005–2015).

We used these susceptibility classifications to trigger further evaluation of potential effects on GDEs. If we determined a GDE unit to have moderate or high susceptibility to changing groundwater conditions, we used biological information to assess whether evidence exists of a biological response to changing groundwater levels or degraded groundwater quality. The biological response analysis was based on changes in Normalized Difference Vegetation Index (NDVI) and Normalized Difference Moisture Index (NDMI) data for individual vegetation polygons within the GDE units (Klausmeyer et al. 2019). The polygons correspond to different GDE mapping units (i.e., different species compositions) and the size of the GDE polygons varied.

NDVI, which estimates vegetation greenness, and NDMI, which estimates vegetation moisture, were generated from surface reflectance corrected multispectral Landsat imagery corresponding to the period of July 9 to September 7 of each year, which represents the period when GDE species are most likely to use groundwater (see Klausmeyer et al. 2019 for further description of methods). Vegetation polygons with higher NDVI values indicate increased density of chlorophyll and photosynthetic capacity in the canopy, an indicator of vigorous, growing vegetation. Similarly, high NDMI values indicate that the vegetation canopy has high water content and is therefore not drought stressed. These indices are both commonly used proxies for vegetation health in analyses of temporal trends in health of groundwater-dependent vegetation (Rouse et al. 1974 and Jiang et al. 2006 as cited in Klausmeyer et al. 2019).

Based on the NDVI and NDMI data, groundwater quality data from wells in or near GDE units in the Fillmore and Piru groundwater basins, and the likely susceptibility of the terrestrial and aquatic species and natural communities in each GDE unit to reported groundwater quality constituents, we found no evidence of a biological response associated with groundwater quality in any of the GDE units. Groundwater quality is therefore not addressed further in the analysis of potential effects.

## 5.2 Biological Data

Tracking the health of all components of groundwater dependent ecosystems through time would involve systematic tracking of populations through time and accounting for changes in driving variables such as floods, climate, and other stressors on populations. This section focuses on changes in vegetation through time using remote sensing data. While increases or decreases in vegetation health do not provide a definitive indication that other components of the ecosystem are thriving or under stress, it provides a reasonable first-order check on the clear linkage between groundwater and the other communities that compose the ecosystem. Previous work has shown that decreases in vegetation vigor are correlated to decreases in remote sensing metrics such as NDVI (e.g., Huntington et al. 2015) and that decreases in vegetation health often corelate with decreases in overall ecosystem health. Tracking the change in NDVI and NDMI for individual polygons shows how the greenness of those polygons change through time. It is crucial to remember that the Santa Clara River and its tributaries in the Fillmore and Piru groundwater basins are dynamic braided rivers that shift through time. This shifting uproots vegetation and creates new surfaces for seedlings to establish. Following floods, the proximity to the river channel (and hence distance to groundwater) as well as the relative elevation of a given vegetation polygon may change. It is therefore useful to average changes over the different GDE units to account for these changes.

#### 5.2.1 Piru Groundwater Basin

## Del Valle Riparian Complex

The mean NDVI and NDMI for the Del Valle Riparian Complex from 1985–2018 were 0.41 and 0.08, respectively. The Del Valle Riparian Complex had relatively steady NDVI values with drops during the early 1990s drought, the 2005 flood, and after 2015. The NDVI values were lower than the other forested riparian complexes (e.g., the East Grove Riparian Complex and the Cienega Riparian Complex), but the declines during the drought were much less severe than for the Cienega Riparian Complex. Mean NDMI has declined slowly since 1995, but the overall change was relatively small. This site has experienced extensive changes in vegetation in response to floods, as visible from comparison of aerial photographs. The Santa Clara River in this GDE unit has a highly dynamic braided channel that is subject to extensive erosion and deposition, even during smaller floods which uprooted vegetation at the site. There is no evidence that the decline in NDVI (which is very small) was due to changes in groundwater management (Figure 5.2-1).

From 2011-2020 NDVI increased as indicated by the slope of a best-fit line to NDVI values through time (Figure 5.2-2). Using the slope of NDVI through time limits the importance of individual years and gives a more representative picture of NDVI changes. For the Del Valley Riparian Complex, NDVI increased from 2011-2020 in the upstream sections of the riparian complex and decreased at the downstream end, likely due to decreased extent of rising groundwater during the drought.





Figure 5.2-1. NDVI and NDMI for the Del Valle Riparian Complex GDE Unit in the Piru Groundwater Basin.



Figure 5.2-2. Slope of NDVI changes from 2011-2020 for the Del Valle Riparian Complex. Blue represents increases in NDVI while red represents decreases in NDVI.

## Piru Basin Santa Clara River Riparian Shrub

The mean NDVI and NDMI for the Santa Clara River Riparian Complex in the Piru Groundwater Basin from 1985–2018 were 0.21 and -0.11, respectively (Figure 5.2-3). These values are relatively low compared to other GDE units in the Fillmore and Piru groundwater basins. The NDVI and NDMI peaked in 1995, 2006, and 2011, following relatively wet years, and declined back to background levels between these peaks. There is no long-term trend in either NDVI or NDMI for this GDE unit (Figure 5.2-3).



**Figure 5.2-3.** NDVI and NDMI for the Santa Clara River Riparian Shrub GDE Unit in the Piru Groundwater Basin.

## Piru Creek Riparian GDE Unit

The mean NDVI and NDMI for the Piru Creek Riparian GDE from 1985—2018 were 0.36 and -0.026, respectively. NDVI values have been relatively steady, with a small decrease in 1996 and a gradual increase starting in 2006 through 2018 (Figure 5.2-4). There was only a small decline in NDVI during the 2012–2016 drought.




Figure 5.2-4. NDVI and NDMI for the Piru Creek Riparian GDE Unit.

## Piru Basin Tributary Riparian GDE

The mean NDVI and NDMI for the Tributary Riparian GDE Unit in the Piru Groundwater Basin from 1985–2018 were 0.30 and -0.07, respectively. The NDVI was relatively steady from 1989–2007 before increasing in 2008 and remaining relatively steady through 2018 (Figure 5.2-5). NDVI was relatively steady during the 1989–1991 and 2012–2016 droughts. Drops in NDVI occurred in 1996 and 2002, but were small compared to changes in other GDE units. NDMI varied more than NDVI from 1989–2007, and had a similar increase in 2008, but declined from 2010–2018 to the mean value over the period of record.





Figure 5.2-5. NDVI and NDMI for the Piru Basin Tributary Riparian GDE Unit.

## 5.2.2 Fillmore Groundwater Basin

## Fillmore Basin Santa Clara River Riparian Shrub

The mean NDVI and NDMI for the Santa Clara River Riparian Shrub GDE in the Fillmore Groundwater Basin from 1985–2018 were 0.29 and -0.046, respectively (Figure 5.2-6). The Santa Clara River Riparian Shrubland in the Fillmore Groundwater Basin has relatively low NDVI values typical of low-density vegetation in the GDE. The NDVI varies over 2- to 5-year cycles (Figure 5.2-6). The NDVI and NDMI values declined during the early 1990s and 2012–2016 drought. NDVI and NDMI have not recovered from the most recent drought (Figure 5.2-6).



**Figure 5.2-6.** NDVI and NDMI for the Santa Clara River Riparian Shrub GDE Unit in the Fillmore Groundwater Basin.

#### Cienega Riparian Complex

The NDVI analysis of the Cienega Riparian Complex focuses on the Fillmore Basin where most of the mapped GDEs occur. The mean NDVI and NDMI values for the Cienega Riparian Complex from 1985–2018 were 0.48 and 0.12, respectively. The mean NDVI in the Cienega Riparian Complex was relatively consistent between 1995 and 2014 (Figure 5.2-7) but dropped by about half during the 1989–1991 and 2012–2016 droughts (Figure 5.2-7). Following the early 1990s drought the NDVI recovered to its pre-drought value within 2 to 3 years. A similar recovery did not occur following the 2012 to 2016 drought and field observation confirm that much of the willow and cottonwood forest died during the drought (Figure 5.2-8). The 2005 flood was the only major flood during this period and resulted in a short-term increase in NDVI, likely because the wet water year supported extensive vegetation growth and new surfaces were rapidly colonized.

NDMI values declined slightly from 1995–2014 and had similar drops during the early 1990s and 2012–2016 droughts. The reason for the decline in NDMI from 1995–2014 is not known. A similar drop in vegetated health using a different remote sensing technique was observed by Kibler et al. (2019). Anecdotal evidence suggests that some of the Arundo stands are recovering following the drought, but much of the willow and cottonwood forest in the GDE died during the drought.




Figure 5.2-7. NDVI and NDMI for the Cienega Riparian Complex GDE Unit.




Figure 5.2-8. Slope of NDVI change in the Cienega Riparian Complex in the Fillmore and Piru Basins from 2011-2020. Red areas have declining NDVI and Blue areas have increasing NDVI.

## East Grove Riparian Complex

The mean NDVI and NDMI for the East Grove Riparian Complex from 1985–2018 were 0.52 and 0.16, respectively. NDVI values for the East Grove Riparian Complex were relatively steady through about 1997, with a small drop following the early 1990s drought (Figure 5.2-9).

Starting around 1998 the NDVI steadily increased until declining in 2013. There was a slight increase in NDVI following the 2005 flood. Mean NDMI was relatively constant until increasing in 2005. NDMI declined more than NDVI during the 2012–2016 drought (Figure 5.2-9).




Figure 5.2-9. NDVI and NDMI for the East Grove Riparian Complex GDE Unit.

Between 2011 and 2020 the slope of the NDVI through time was positive (i.e., NDVI increased), with decreases where the channel shifted (Figure 5.2-10). Decreases in NDVI occurred at the upstream portion of the East Grove Complex, reflecting a change in the flow path of the Santa Clara River as well as a decline in vegetation health on the southeast portion of the GDE. In comparison with the Cienega Riparian Complex, the increasing NDVI in the East Grove Complex suggests that groundwater levels did not drop below the rooting zone of the riparian complex and hence the GDE was much more resilient.




Figure 5.2-10. Slope of NDVI change in the Cienega Riparian Complex in the Fillmore and Piru Basins from 2011-2020. Red areas have declining NDVI and Blue areas have increasing NDVI.

#### Fillmore Basin Tributary Riparian GDE Unit

The mean NDVI and NDMI for the Tributary Riparian GDE in the Fillmore Groundwater Basin from 1985–2018 were 0.50 and 0.058, respectively (Figure 5.2-11). The NDVI was relatively steady through the 1989-1991 drought and through 2017 before dropping in 2018 likely due to vegetation mortality due to the Thomas Fire. Short-term drops in NDVI occurred in 1996 and 2002, but were small compared to changes in other GDE units. NDMI declined in 2012 at the start of the 2012–2016 drought, and has remained below the mean. In 2018 the NDMI further declined due to the Thomas Fire (Figure 5.2-11).




Figure 5.2-11. NDVI and NDMI for the Fillmore Basin Tributary Riparian GDE Unit.

## Sespe Creek Riparian

The mean NDVI and NDMI for the Sespe Creek Riparian Complex from 1985–2018 were 0.39 and 0.084, respectively. The Sespe Creek Riparian GDE has relatively steady intermediate NDVI of 0.3–0.4 prior to 2005, at which point NDVI dropped (Figure 5.2-12) as a result of the flood. Following the 2005 flood, NDVI gradually increased, as would be expected if the riparian forest was recovering following the flood. NDVI dropped from 2015–2018 (Figure 5.2-12). NDMI has been relatively consistent through time (Figure 5.2-12).




Figure 5.2-12. NDVI and NDMI for the Sespe Creek Riparian Complex GDE Unit.

## 5.3 Climate Change Effects

The effects of climate change on groundwater and interconnected surface water are discussed in DBS&A 2021. As an overview, the future groundwater levels forecast with assumed climate change factors (2070CF [climate change factor]) are not materially different from those recorded during the historical record. There is no suggestion of long-term chronic declines in groundwater levels.

Low water levels associated with major droughts (and accounting for future climate) are lower (typically 10–30 ft) than those of the historical time period (i.e., without the effects of climate change). Despite the lower, drought-induced water levels, the water levels return to historical high water level conditions during subsequent wet to normal precipitation periods. Statewide climate models suggest that there could be an increase in the duration, severity, and frequency of droughts and extreme floods through the remainder of 21<sup>st</sup> century (Swain et al. 2018) could impact groundwater availability to GDEs but may also increase the frequency of flood events that are crucial for regrowth of the pioneer species that make up the GDEs along the Santa Clara River.

Climate change may alter the water demands of groundwater dependent vegetation, but the response is complex because decreased transpiration associated with increased carbon dioxide in

the atmosphere may counter increased evaporation due to temperature increases (e.g., Klove et al. 2014).

## 5.4 Summary of Potential Effects

Potential effects on each GDE unit are summarized here based on three primary criteria:

- 1. Ecological value (high, moderate, low), as described in Section 4.3.
- 2. Ecological condition of the GDEs within each unit (good, fair, poor), based on the information summarized in Section 4.1 and the NDVI/NDMI data presented in Section 5.2.
- 3. Susceptibility to changing groundwater conditions (high, moderate, low) based on available hydrologic data, climate change projections, and the GDE susceptibility classifications summarized in Table 5.1-1.

The groundwater dependence of each GDE unit, including any rising groundwater, is also summarized to provide context for the effects assessment.

#### 5.4.1 Piru

#### Del Valle Riparian Complex GDE Unit

Groundwater Dependence: Likely

- Shallow groundwater measurements are rare in this unit but the historical persistence of the riparian forest and widespread willows and cottonwoods suggest that groundwater is likely within the rooting zone of plants.
- This GDE Unit is a mixture of willows and cottonwoods that are likely connected to groundwater and facultative phreatophytes (e.g., mulefat and arrow-weed thickets) that may be connected to groundwater.
- Perennial surface water flows are likely connected with groundwater.

#### Ecological Value: High

- The Del Valle Riparian Complex GDE Unit supports a relatively large number of specialstatus species and ecological communities, some of which are directly dependent on groundwater.
- The Unit includes designated critical habitat for two federally listed species.
- The Unit supports species and ecological communities that are vulnerable to changes in groundwater levels.

#### Ecological Condition: **Good**

• NDVI/ NDMI from 1985–2018 was relatively constant until a flood-related reduction in NDVI in 2005 followed by a gradual recovery until 2015, at which point NDVI began a gradual decline. As of 2018, NDVI had not declined below the long-term average. NDMI has declined slowly since 1995 and in 2018 was below the long-term average. These trends suggest that the structure and function of riparian vegetation in the Unit may have experienced declines during the drought, particularly at its downstream end. Although groundwater levels are relatively stable, and this reach receives wastewater effluent from upstream, the limited well data in this GDE unit are insufficient to determine whether

- groundwater pumping has contributed to the observed declining vegetation condition at the downstream end of the unit.
- Habitat suitability in the downstream portion of the Unit may be somewhat compromised by the decline in vegetation condition for special-status species that rely on vegetation (e.g., riparian birds).
- Groundwater contributes to the ecological function and habitat value of the Santa Clara River, which supports native aquatic and semi-aquatic species and beneficial uses in and adjacent to the Unit.

#### Susceptibility to Changing Groundwater Conditions: Moderate

- Shallow groundwater conditions in the Unit since the 1930s have fluctuated considerably in response to drought and possibly other factors. An 80-ft decline with the 2012–2016 drought has been followed by a slow recovery, and by spring 2020 groundwater depth was again within the baseline range for non-drought conditions. The groundwater well is at the downstream end of the unit and therefore may not reflect groundwater elevation changes further upstream.
- Rooting depths of willows and cottonwoods in this unit range up to 6.9 ft while the average relative elevation is 10 ft for cottonwoods and 5.6 ft for red willow. The mulefat thickets are not likely connected to groundwater.
- The decline in groundwater at the downstream end of the unit exceeded the rooting depth of the vegetation. Groundwater declines in the rest of the unit were likely less based on the vegetation response to the drought, but there are no well data further upstream.
- Future changes in groundwater conditions in the Unit related to increased groundwater pumping, declining effluent releases from upstream, or climate change could cause groundwater levels to fall below the baseline range and result in mortality of the trees that comprise the GDE. Projections of climate change and groundwater pumping in the future suggest that changes in groundwater elevation are unlikely.

The Unit includes a perennial portion of the mainstem Santa Clara River that is considered an interconnected surface water. The degree to which interconnected surface waters in this reach are maintained by releases from upstream effluent sources is unknown, but is believed to be significant.

#### Potential for effects

Available data are insufficient to discern a clear effect on GDEs related to groundwater management in the Del Valle Riparian Complex GDE Unit. Declines in vegetation health (as shown by NDVI decreases) at the downstream end of the reach suggest that GDEs in this unit are susceptible to drought conditions. However, the extent of GDEs in the unit is moderately susceptible to future decrease in groundwater elevation and surface water conditions and the synergistic effects of climate change (described in Section 5.3). In combination these changes could cause groundwater levels to fall below the baseline range and result in mortality of the trees that comprise the GDE and reduce the extent of the GDE. GDEs in this unit are not expected to experience future water levels that are lower than the historical period, but more frequent or longer duration droughts due to climate change could also affect the extent of vegetation-dominated GDEs.

Monitoring of ecological conditions and trends in vegetation-dominated GDEs and in the Santa Clara River using remote sensing is recommended and is further discussed in Section 6 GDE

Monitoring. Changes to upstream effluent releases may impact aquatic habitat and groundwater elevation in this GDE unit.

#### Santa Clara River Riparian Shrubland GDE Unit

## Groundwater Dependence: Possible

- There are few shallow groundwater measurements in this unit. Spring 2019 water contours provided by United water showed groundwater levels within 5-10 of the ground surface in parts of the unit.
- This GDE unit includes a large polygon of giant reed (Arundo) at the downstream end of Piru Groundwater Basin, with patches of sandbar willows and large mulefat thickets. Given the shallow rooting depth of mulefat thickets (approximately 2 ft), they likely are not connected to groundwater. Other vegetation communities in the unit may be connected to groundwater. Small patches of sandbar willows are present in this unit and have average relative elevations in the Santa Clara River of 4.8 ft and the relative elevation ranges up to 9 ft (Appendix C).
- Ephemeral surface water flows are not connected with groundwater.

#### Ecological Value: Moderate

- The Santa Clara River Riparian Shrubland GDE Unit supports a moderate number of special-status species and ecological communities, some of which are directly dependent on groundwater.
- The Unit includes designated critical habitat for two federally listed species.
- This naturally intermittent reach supports regionally rare alluvial scrub habitat.
- The Unit supports species and ecological communities that are somewhat vulnerable to changes in groundwater levels. Although the Santa Clara River in the Unit provides migration habitat for Southern California steelhead and Pacific lamprey, the migration habitat has low vulnerability to groundwater reduction because most fish migration occurs during seasonal high surface water flow periods.

#### Ecological Condition: Fair

- NDVI/ NDMI values in the Unit from 1985–2018 are low compared to other GDE units in
  the Fillmore and Piru groundwater basins, reflecting the relatively sparse vegetation. Mean
  NDVI and NDMI during this period increased in response to wet years and returned to
  long-term average values between the peaks. Mean NDVI or NDMI values in the Unit do
  not appear responsive to drought. There is no evidence that groundwater management
  affects the ecological condition of this GDE unit.
- Current habitat suitability for those special-status species with likelihood to occur in the Unit may have declined relative to historical conditions.
- Groundwater provides little or no contribution to the ecological function and habitat value of the Santa Clara River in the Unit, which is intermittent and mainly supports seasonal migration habitat for anadromous fishes.

#### Susceptibility to Changing Groundwater Conditions: Low

• Shallow groundwater conditions in the Unit since the 1970s have fluctuated in response to drought, with sharp drops followed by recovery to pre-drought levels. As of fall 2019, the shallow groundwater level recorded in the Unit had apparently recovered from the large

drop associated with the 2012–2016 drought and was again within the baseline range. So long as the duration and frequency of droughts does not change, the effects on this GDE unit are expected to be minimal. The sandbar willows and eucalyptus occur on the margin of the unit adjacent to agricultural lands and may subsist on agricultural runoff.

- Future changes in groundwater conditions in the Unit related to increased groundwater production or climate change are not expected to cause groundwater levels to fall below the baseline range. As a result the potential effects on GDEs are deemed negligible.
- The Unit includes an intermittent reach of the mainstem Santa Clara River that does not provide perennial aquatic habitat or beneficial uses.

#### Potential for effects

Modeling suggests that groundwater levels are likely to be stable in this reach. Moreover, the vegetation that makes up this unit may use groundwater when groundwater levels are high in the spring, but high groundwater levels are likely not persistent in this unit. The unit is therefore likely not strongly dependent upon groundwater and is comprised of sparse low water use species with relatively shallow rooting depths. Therefore, the potential for effects on this unit is low.

Monitoring of ecological conditions and trends in vegetation-dominated GDEs and in the Santa Clara River using remote sensing is recommended and is further discussed in Section 6 GDE Monitoring.

#### Cienega Riparian Complex GDE Unit

Groundwater Dependence: Certain

• Rising groundwater in this unit provides surface flows and keeps groundwater within the rooting zone (5–15 ft) of the vegetation.

## Ecological Value: High

- The Cienega Riparian Complex GDE Unit supports a moderate number of special-status species, some of which are directly dependent on groundwater.
- The Unit includes designated critical habitat for two federally listed species.
- The Unit supports species and ecological communities that are vulnerable to changes in groundwater levels.

#### Ecological Condition: Fair

- NDVI/ NDMI trends from 1985–2018 indicate vegetation responds mainly to precipitation and runoff (e.g., drought) but a decline in NDMI from 2005–2014 suggests other factors, potentially including groundwater management, may influence vegetation condition in the Unit. As of 2018 the vegetation structure and functions in the Cienega Riparian Complex GDE unit are no longer intact or within the range of natural variability, due at least in part to vegetation mortality from the 2012–2016 drought. The degree to which groundwater management has exacerbated the decline in groundwater elevation during the recent and other droughts is not known.
- Habitat suitability in the Unit for special-status species relying on vegetation (e.g., riparian birds)may be compromised by the decline in vegetation condition during droughts.
- Groundwater contributes to the ecological function and habitat value of the Santa Clara River, which supports native aquatic and semi-aquatic species and beneficial uses in and adjacent to the Unit.

#### Susceptibility to Changing Groundwater Conditions: High

- Shallow groundwater conditions in the Unit since the 1970s have fluctuated in response to drought, with sharp drops followed by recovery within about one year. By 2019–2020, shallow groundwater levels recorded at the single well in the Unit were again within the baseline range for the period of record, but the native cottonwoods and willows died during the drought and have not yet recovered. Those species are anticipated to recover once a future flooding event(s) removes the debris and the land surface is but conditioned for repopulation. The time required for that recovery is unknown.
- Reported maximum rooting depths of willows and cottonwoods in this unit range up to 6.9
  ft while the average relative elevation is 10 ft for cottonwoods and 5.6 ft for red willow.
  The mulefat thickets are not likely connected to groundwater.
- Future changes in groundwater conditions in the Unit related to increased groundwater production or climate change could cause groundwater levels to fall below the baseline range and result in potential mortality to vegetation that comprises the GDE. Projections of climate change and groundwater pumping in the future suggest that changes in groundwater elevation are unlikely. However, based on widespread tree mortality during the 2012–2016 drought, future changes in the frequency or duration of droughts similar to 2012–2016 could have a deleterious effect on the GDE.
- The Unit includes a perennial portion of the mainstem Santa Clara River, which is considered an interconnected surface water.

#### Potential for effects

Modeling suggests that climate change is unlikely to cause groundwater levels to drop below the baseline range. However, changes to the duration or severity of droughts could impact the health of the GDE through increased tree mortality. Moreover, it is possible that arundo could replace the cottonwood and willow forests that died during the 2012–2016 drought which would lead to a decrease in habitat for other species (i.e., riparian birds).

Monitoring of ecological conditions and trends in vegetation-dominated GDEs and in the Santa Clara River using remote sensing is recommended and is further discussed in Section 6 GDE Monitoring. For this unit, coupling remote sensing and shallow groundwater elevation monitoring, particularly during and following droughts, is recommended.

## Piru Creek Riparian GDE Unit

Groundwater Dependence: Uncertain/Unlikely

- F
- Groundwater wells in the rooting zone of plants (<30 ft) are rare in this unit.
- Releases from Santa Felicia Dam sustain surface flows.

#### Ecological Value: High

- The Piru Creek Riparian Complex GDE Unit supports a relatively high number of specialstatus species and ecological communities, some of which are directly dependent on groundwater.
- The Unit includes designated critical habitat for two federally listed species.
- The Unit supports species and ecological communities whose habitat in the Unit may be vulnerable to changes in groundwater levels.

## Ecological Condition: Good

- NDVI/ NDMI from 1985–2018 was relatively constant and seemingly unresponsive to droughts and floods. These trends suggest that the structure and function of riparian vegetation in the Unit are relatively intact and within the range of natural variability. Riparian vegetation in the Unit may be sustained by releases from Santa Felicia Dam which likely raise the groundwater level in this area. Available information indicates that adverse impacts are not likely occurring in the Unit, at least partially as a result of current surface water releases that provide water to at least the near-channel portions of the GDE unit even if groundwater is below the rooting depth of most riparian plants.
- Suitable habitat is present for those special-status species with likelihood to occur in the Unit.
- Releases from Santa Felicia Dam likely raise groundwater levels and help maintain baseflows in Piru Creek, thus contributing to the ecological function and habitat value of Piru Creek. Piru Creek supports native aquatic and semi-aquatic species and beneficial uses in and adjacent to the Unit.

## Susceptibility to Changing Groundwater Conditions: Undetermined, likely low

- There are no monitoring wells in the unit and shallow groundwater conditions and trends in the Unit are therefore unknown.
- Assuming continued releases from Santa Felicia Dam, it is unlikely that future changes in
  groundwater conditions in the Unit related to increased groundwater production or climate
  change will cause groundwater levels to fall below the baseline range. As a result the
  potential effects on GDEs are deemed negligible.
- Piru Creek in this GDE unit has perennial flow due to releases from Santa Felicia Dam, but surface flow is not connected to groundwater. The lower portion of Piru Creek near the confluence with the Santa Clara River periodically lacks surface flow. As described previously, releases from Santa Felicia Dam likely raise groundwater levels and help maintain baseflows in Piru Creek.

#### Potential for effects

Available data are insufficient to discern a clear effect on GDEs related to groundwater management in the Piru Creek Riparian Complex GDE Unit. However, groundwater levels and baseflows in Piru Creek are likely maintained by releases from Santa Felicia Dam, thus the susceptibility of GDEs in the Unit (i.e., vegetation mortality) to future changes in groundwater conditions and the synergistic effects of climate change is low. With continued dam releases, the potential for these combined effects to cause groundwater levels to fall below the baseline range and result in potential effects on GDEs is low.

Monitoring of ecological conditions and trends in vegetation-dominated GDEs and in Piru Creek using remote sensing is recommended and is further discussed in Section 6. Coupling remote sensing with a shallow groundwater monitoring well would help to assess the degree to which groundwater dynamics affect GDEs in this unit. In this GDE unit, changes to releases from Santa Felicia Dam may affect aquatic habitat and groundwater elevation.

#### Tributary Riparian GDE Unit

Groundwater Dependence: Unlikely

• There are no shallow groundwater measurements in this unit. Based on the position in the landscape a connection to the regional aquifer is unlikely.

- This GDE unit contains a mixture of obligate and facultative phreatophytes that may be connected to groundwater (unlikely) or surviving on episodic surface water flows.
- Ephemeral surface water flows are not connected with groundwater.

### Ecological Value: Moderate

- The Tributary Riparian GDE Unit supports a relatively low number of special-status species and ecological communities and the dependence of these species and communities on groundwater is uncertain.
- The Unit includes designated critical habitat for one federally listed species.
- The species and ecological communities in the Unit have low vulnerability to changes in groundwater levels. The tributary streams in this GDE Unit are considered ephemeral and are not connected to groundwater, thus they provide little habitat value for fish and other aquatic species. They do, however, support valuable riparian habitat and likely movement corridors for a variety of native wildlife species.

## Ecological Condition: Fair

- NDVI/ NDMI trends in the Unit from 1985–2018 show relatively little change in vegetation condition during most of this period, with little change in response to droughts or floods. It is unlikely that adverse impacts are occurring in the Unit as a result of current groundwater management.
- Suitable habitat is present for those special-status species with likelihood to occur in the Unit.
- Groundwater likely provides little or no contribution to the ecological function and habitat
  value of the ephemeral tributaries in the Unit, which support vegetation but have little
  habitat value for fish or other aquatic species.

## Susceptibility to Changing Groundwater Conditions: Moderate

- There are no monitoring wells in the unit and shallow groundwater conditions and trends in the Unit are therefore unknown.
- Model results suggest that the groundwater elevations are unlikely to decline under climate change. Moreover, because this GDE unit is disconnected from the aquifer, future changes in groundwater conditions in the Unit related to increased groundwater production or climate change are unlikely to cause groundwater levels to fall below the baseline range and result in mortality to vegetation that comprises the GDE.
- Streams within the Unit includes are ephemeral and do not provide perennial aquatic habitat or beneficial uses.

## Potential for effects

Based on the position of this GDE unit in the watershed it is unlikely that groundwater management will affect the health of the GDE. Model results suggest that the groundwater levels will remain constant in the Fillmore and Piru Basins under climate change (DBS&A 2021). If groundwater pumping were to increase in this GDE unit, monitoring of groundwater levels and GDE health (using remote sensing) would be necessary. GDEs in the unit likely have low susceptibility to future changes in groundwater conditions and the synergistic effects of climate change.

#### 5.4.2 Fillmore

## Santa Clara River Riparian Shrubland GDE Unit

## Groundwater Dependence: Possible

- There are few shallow groundwater measurements in this unit. Spring 2019 water contours provided by United water showed groundwater levels within 5-10 feet of the ground surface in parts of the unit.
- The Santa Clara River Riparian Shrubland GDE unit is primarily made up of vegetation that may or may not rely on groundwater.
- Surface water flows are not interconnected with groundwater.

#### Ecological Value: Moderate

- The Santa Clara River Riparian Shrubland GDE Unit supports a relatively large number of special-status species and ecological communities, some of which are directly dependent on groundwater.
- The Unit includes designated critical habitat for two federally listed species.
- This naturally intermittent reach supports regionally rare alluvial scrub habitat.
- The Unit supports species and ecological communities that are vulnerable to changes in groundwater levels. Although the Santa Clara River in the Unit provides migration habitat for Southern California steelhead and Pacific lamprey, the migration habitat has low vulnerability to groundwater reduction because most fish migration occurs during seasonal high surface water flow periods.

#### Ecological Condition: Fair

- NDVI/ NDMI trends in the Unit from 1985–2018 indicate a decline in vegetation condition since 2012 relative to the long-term average, likely in response to decreased precipitation and runoff (e.g., drought). The current vegetation structure and functions may be compromised and somewhat below the range of natural variability. Groundwater management is unlikely to have an adverse effect on this GDE Unit because summer groundwater is typically deeper than the rooting depth of vegetation in the reach.
- Current habitat suitability for those special-status species with likelihood to occur in the Unit may have declined relative to historical conditions.
- Because surface water in this reach is largely disconnected from groundwater, groundwater
  provides little or no contribution to the ecological function and habitat value of the Santa
  Clara River in the Unit, which is intermittent and mainly supports seasonal migration
  habitat for anadromous fishes.

## Susceptibility to Changing Groundwater Conditions: Moderate

- Since 2015, shallow groundwater conditions in the Unit have fluctuated in response to drought, with a sharp drop in 2013 followed by slow recovery. By 2019, the shallow groundwater level recorded in the Unit had nearly returned to the long-term average (i.e., just below the baseline range).
- Future changes in groundwater conditions in the Unit related to increased groundwater
  production or climate change could cause groundwater levels to fall below the baseline
  range and result in mortality to vegetation that comprises the GDE. Projections of climate
  change and groundwater pumping in the future suggest that changes in groundwater
  elevation are unlikely. However, based on widespread tree mortality during the 2012–2016

drought, future changes in the frequency or duration of droughts similar to 2012–2016 could have a deleterious effect on the GDE, particularly at the downstream margin of the unit.

• The Unit includes an intermittent reach of the mainstem Santa Clara River that does not provide perennial aquatic habitat or beneficial uses.

#### Potential for effects

Modeling suggests that groundwater levels near the Santa Clara River Riparian Shrubland GDE unit are unlikely to change due to climate change or modest changes to groundwater pumping. However, GDEs in the Unit are moderately susceptible to future changes in groundwater conditions and the synergistic effects of climate change, which in combination could cause groundwater levels to fall below the baseline range and result in potential effects on GDEs.

Monitoring of ecological conditions and trends in vegetation-dominated GDEs and in the Santa Clara River using remote sensing is recommended and is further discussed in Section 6 GDE Monitoring.

#### East Grove Riparian Complex GDE Unit

Groundwater Dependence: Certain

- This GDE Unit occurs at a site of rising groundwater.
- This GDE Unit is primarily made up of cottonwoods and willows that rely on shallow groundwater.
- Perennial surface water flows are rising groundwater.

## Ecological Value: High

- The East Grove Riparian Complex GDE Unit supports a relatively large number of specialstatus species and ecological communities, some of which are directly dependent on groundwater.
- The Unit includes designated critical habitat for two federally listed species.
- The Unit supports species and ecological communities that are vulnerable to changes in groundwater levels.

#### Ecological Condition: Good

- NDVI/ NDMI trends from 1985–2018 show minimal change in the Unit and indicate
  vegetation responds mainly to precipitation and runoff (e.g., drought). The vegetation
  structure and functions are relatively intact and within the range of natural variability, and
  adverse impacts are not likely occurring in the Unit as a result of current groundwater
  management.
- Suitable habitat is present for those special-status species with likelihood to occur in the Unit.
- With the exception of flow during storm events, Santa Clara River surface flows in this unit are composed of rising groundwater. As a result, groundwater contributes to the ecological function and habitat value of the Santa Clara River, which supports native aquatic and semi-aquatic species and beneficial uses in and adjacent to the Unit.

Susceptibility to Changing Groundwater Conditions: Moderate

- Shallow groundwater conditions in the Unit since the 1960s have fluctuated considerably in response to drought and possibly other factors. By 2019–2020, shallow groundwater levels recorded at the two wells in the Unit were again within the baseline range for each well.
- Recorded maximum rooting depths of willows and cottonwoods in this unit range up to 6.9 ft while the average relative elevation is 10 ft for cottonwoods and 5.6 ft for red willow. Mulefat has a rooting depth of 2 ft (Appendix C), and the mulefat thickets are not likely connected to groundwater.
- Future changes in groundwater conditions in the Unit related to increased groundwater production or climate change could cause groundwater levels to fall below the baseline range and result in mortality to vegetation that comprises the GDE. Projections of climate change and groundwater pumping in the future suggest that changes in groundwater elevation are unlikely. However, because the extent of rising groundwater decreased and vegetation health declined at the upstream end of the unit during the 2012–2016 drought, changes in the frequency or duration of droughts to make 2012-2016 conditions more common could have a deleterious effect on the GDE.
- The Unit includes a perennial portion of the mainstem Santa Clara River, which is an area of rising groundwater.

#### Potential for effects

Modeling suggests that groundwater levels are unlikely to drop below the baseline range due to climate change. However, changes to the duration or severity of droughts could impact the health of the GDE through increased tree mortality.

Monitoring of ecological conditions and trends in vegetation-dominated GDEs and in the Santa Clara River using remote sensing is recommended and is further discussed in Section 6 GDE Monitoring. Additionally, monitoring of surface water flows or groundwater elevations should be conducted to assess any changes to hydrology that might impact GDEs.

#### Cienega Riparian Complex GDE Unit

Groundwater Dependence: Certain

• Rising groundwater in this unit provides surface flows and keeps groundwater within the rooting zone (5-15 ft) of the vegetation.

## Ecological Value: High

- The Cienega Riparian Complex GDE Unit supports a moderate number of special-status species, some of which are directly dependent on groundwater.
- The Unit includes designated critical habitat for two federally listed species.
- The Unit supports species and ecological communities that are vulnerable to changes in groundwater levels.

# Ecological Condition: F

NDVI/ NDMI trends from 1985–2018 indicate vegetation responds mainly to precipitation
and runoff (e.g., drought) but a decline in NDMI from 2005–2014 suggests other factors,
potentially including groundwater management, may influence vegetation condition in the
Unit. As of 2018 the vegetation structure and functions in the Cienega Riparian Complex
GDE unit are no longer intact or within the range of natural variability, due at least in part

to vegetation mortality from the 2012–2016 drought. The degree to which groundwater management has exacerbated the decline in groundwater elevation during the recent and other droughts is not known.

- Habitat suitability in the Unit may be compromised by the decline in vegetation condition during droughts for special-status species relying on vegetation (e.g., riparian birds).
- Groundwater contributes to the ecological function and habitat value of the Santa Clara River, which supports native aquatic and semi-aquatic species and beneficial uses in and adjacent to the Unit.

#### Susceptibility to Changing Groundwater Conditions: High

- Shallow groundwater conditions in the Unit have fluctuated in response to drought since the 1970s, with sharp drops followed by recovery within about one year. By 2019–2020, shallow groundwater levels recorded at the single well in the Unit were again within the baseline range for the period of record, but the native cottonwoods and willows died during the drought and have not yet recovered. Those species are anticipated to recovery once a future flooding event(s) removes the debris and the land surface is better conditioned for repopulation. The time required for that recovery is unknown.
- Reported maximum rooting depths of willows and cottonwoods in this unit range up to 6.9 ft while the average relative elevation is 10 ft for cottonwoods and 5.6 ft for red willow. The mulefat thickets are not likely connected to groundwater.
- Future changes in groundwater conditions in the Unit related to increased groundwater production or climate change could cause groundwater levels to fall below the baseline range and result in potential mortality to vegetation that comprises the GDE. Projections of climate change and groundwater pumping in the future suggest that changes in groundwater elevation are unlikely. However, based on widespread tree mortality during the 2012–2016 drought, future changes in the frequency or duration of droughts similar to 2012–2016 could have a deleterious effect on the GDE.
- The Unit includes a perennial portion of the mainstem Santa Clara River, which is considered an interconnected surface water.

#### Potential for effects

Modeling suggests that groundwater levels are unlikely to drop below the baseline ranged due to climate change. However, changes to the duration or severity of droughts could impact the health of the GDE through increased tree mortality. Moreover, it is possible that arundo could replace the cottonwood and willow forests that died during the 2012 2016 drought which would lead to a decrease in habitat for other species (i.e., riparian birds).

Monitoring of ecological conditions and trends in vegetation-dominated GDEs and in the Santa Clara River using remote sensing is recommended and is further discussed in Section 6 GDE Monitoring. For this unit, coupling remote sensing and shallow groundwater elevation monitoring, particularly during and following droughts, is recommended.

## Tributary Riparian GDE Unit

## Groundwater Dependence: Uncertain

• There are no shallow groundwater measurements in this unit. Based on the position in the landscape a connection to the regional aquifer is unlikely.

- This GDE unit is primarily made up of coast live oaks which are facultative phreatophytes that may be connected to groundwater or episodic surface water flows associated with storm events.
- Ephemeral surface water flows are not connected with groundwater.

#### Ecological Value: Moderate

- The Tributary Riparian GDE Unit supports a relatively low number of special-status species and ecological communities and the dependence of these species and communities on groundwater is uncertain.
- There is no designated critical habitat in the Unit.
- The species and ecological communities in the Unit have low vulnerability to changes in groundwater levels. The tributary streams in this GDE Unit are considered ephemeral and are not connected to groundwater, thus they provide little habitat value for fish and other aquatic species. They do, however, support valuable riparian habitat and likely movement corridors for a variety of native wildlife species.

## Ecological Condition: Fair

- NDVI/NDMI trends in the Unit from 1985–2018 indicate relatively little change in vegetation condition during most of this period, with a decline in response the most recent drought (2012–2016) and potential damage to vegetation in 2018 resulting from the Thomas Fire. As a result, the current vegetation structure and functions may be compromised and below the range of natural variability, but it is unlikely that adverse impacts are occurring in the Unit as a result of current groundwater management.
- Current habitat suitability for those special-status species with likelihood to occur in the Unit may have declined relative to historical conditions.
- Groundwater provides little or no contribution to the ecological function and habitat value of the ephemeral tributaries in the Unit, which support vegetation but have little habitat value for fish or other aquatic species.

## Susceptibility to Changing Groundwater Conditions: Low

- There are no monitoring wells in the unit and shallow groundwater conditions and trends in the Unit are therefore unknown, but the position of the tributaries suggests they are unlikely to be linked to regional groundwater.
- Because this GDE unit is disconnected from the aquifer, future changes in groundwater
  conditions in the Unit related to increased groundwater production or climate change are
  unlikely to cause groundwater levels to fall below the baseline range and result in mortality
  to vegetation that comprises the GDE.
- Streams within the Unit includes are ephemeral and do not provide perennial aquatic habitat or beneficial uses.

#### Potential for effects

Based on the position of this GDE unit in the watershed it is unlikely that groundwater management will affect the health of the GDE. If groundwater pumping were to increase in this GDE unit monitoring of groundwater levels and GDE health (using remote sensing) would be necessary. GDEs in the Unit likely have low susceptibility to future changes in groundwater conditions and the synergistic effects of climate change.

### Sespe Creek Riparian Complex GDE Unit

#### Groundwater Dependence: Possible

- This GDE Unit occurs downstream of the confined canyon reach of Sespe Creek.
- Water levels in groundwater are typically > 30 ft, but there are few wells within the shallow groundwater zone.
- This GDE Unit is primarily made up of willows and cottonwoods that rely on shallow groundwater or surface water and some communities (e.g., mulefat) that may rely on groundwater for part of their water needs.
- Surface water flows are perennial for the upper portions of the reach and intermittent downstream. The connection to groundwater in the upper portion is unknown but unlikely.

#### Ecological Value: Moderately High

- The Sespe Creek Riparian Complex GDE Unit supports a moderate number of specialstatus species and ecological communities, some of which are directly dependent on groundwater.
- The Unit includes designated critical habitat for one federally listed species.
- The Unit supports species and ecological communities whose habitat in the Unit may be vulnerable to changes in groundwater levels.

## Ecological Condition: Good

- NDVI/ NDMI from 1985–2018 was relatively constant until a sharp flood-related reduction in NDVI in 2005 followed by a gradual recovery until 2015, at which point NDVI began a gradual decline. As of 2018, NDVI was still within the historical range of variability. NDMI has fluctuated little during the period of record. These trends suggest that the structure and function of riparian vegetation in the Unit are relatively intact and within the range of natural variability. Available information indicates that adverse impacts are not likely occurring in the Unit as a result of current groundwater management. Invasive species, particularly Arundo, are a continuing threat to existing GDEs in this unit.
- Suitable habitat is present for those special-status species with likelihood to occur in the Unit.
- It is undetermined if or to what extent groundwater contributes to the ecological function and habitat value of Sespe Creek, which supports native aquatic and semi-aquatic species and beneficial uses in and adjacent to the Unit.

# Susceptibility to Changing Groundwater Conditions: Lo

- There are no shallow monitoring wells in the unit and shallow groundwater conditions and trends in the Unit are therefore unknown.
- Modeling suggests that groundwater elevations are unlikely to change due to changes in climate or groundwater pumping in the future. Changes to the duration or extent of droughts may cause tree mortality within the GDE unit.
- The upper 2 miles or so of Sespe Creek in this GDE unit are considered perennial, while the lower portion of Sespe Creek is likely intermittent. Sespe Creek's connection to groundwater is undetermined.

#### Potential for effects

Modeling suggests that groundwater elevations are unlikely to change due to changes in climate or groundwater pumping in the future. However, changes in the duration or severity of droughts could impact the health of the GDE through increased physiological stress to riparian vegetationt lealding to branch dieback or whole tree mortality.

Monitoring of ecological conditions and trends in vegetation-dominated GDEs and in Sespe Creek using remote sensing is recommended and is further discussed in Section 6 GDE Monitoring. Additionally, monitoring of surface water flows or groundwater elevations should be conducted to assess any changes to hydrology that might impact GDEs. Further assessing the extent of interconnected surface water in the reach will help to determine potential groundwater impacts to aquatic habitat.

# 5.4.3 GDEs Important for GSP A system

The evaluations of the GDE units in the Fillmore and Piru basins suggests that the following units are the most important for inclusion in the GSP analyses and the development of Sustainable Management Criteria:

- Del Valle Riparian Complex;
- Cienega Riparian Complex; and
- East Grove Riparian Complex.

These units encompass areas of rising groundwater and have hytorically supported large, teeforested wetland complexes.

## 6 GDE MONITORING

GDEs were considered as part of the groundwater Monitoring Program (Appendix XX). Remote sensing of GDEs using NDVI or other widely available data is recommended to monitor the health of GDEs through time. It is expected that NDVI will exhibit some natural variability given the dynamic nature of this river system. For example, NDVI would be expected to initially decline following the large floods that tend to uproot vegetation and provide fresh bare mineral surfaces and appropriate hydrological conditions for seedling establishment of cottonwoods and willows, but then increase again as native cottonwood and willow vegetation becomes established and individual shrubs and trees develop and mature. Groundwater wells near the basin boundaries (the Del Valle, Cienega, and East Grove riparian complexes) can be used to determine changes in groundwater levels. Such data on groundwater levels through time could then be examined to see if there are clear correlations with observed trends in NDVI or related indicators of GDE health.

## 7 PROJECTS AND MANAGEMENT ACTIONS

Projects and management actions (PMAs) are discussed in Section 4 of the GSP. At time of the release of this technical memorandum, the FPBGSA had not determined that projects and/or management actions were needed to sustainably manage the groundwater resources in the Fillmore or Piru basins.

## 8 LITERATURE CITED

ACS (Aquatic Consulting Services, Inc.). 2002. Aquatic Surveys Along the Santa Clara River Part III: West of Commerce Center Bridge to the Ventura County Line, California. Prepared for: The Newhall Land and Farming Company Valencia, California.

Bell, G. P. 1994. Biology and growth habits of giant reed (*Arundo donax*). Pages 1-6 in N. E. Jackson, P. Frandsen, and S. Douthit, editors. *Arundo donax* Workshop Proceedings, November 19, 1993, Ontario, California.

Bell, G. P. 1997. Ecology and Management of *Arundo donax*, and Approaches to Riparian Habitat Restoration in Southern California. Pages 103-113 in Brock, J. H., Wade, M., Pysek, P., and Green, D. (Eds.). Plant Invasions: Studies from North America and Europe. Blackhuys Publishers, Leiden, The Netherlands.

Beller, E. E., P. W. Downs, R. M. Grossinger, B. K. Orr, and M. N. Soloman. 2016. From past patterns to future potential: using historical ecology to inform river restoration on an intermittent California river. Landscape Ecology, 31:581-600, DOI 10.1007/s10980-015-0264-7

Beller, E. E., R. M. Grossinger, M. N. Salomon, S. J. Dark, E. D. Stein, B. K. Orr, P. W. Downs, T. R. Longcore, G. C. Coffman, A. A. Whipple, R. A. Askevold, B. Stanford, and J. R. Beagle. 2011. Historical ecology of the lower Santa Clara River, Ventura River, and Oxnard Plain: an analysis of terrestrial, riverine, and coastal habitats. Prepared for the State Coastal Conservancy. A report of SFEI's Historical Ecology Program, SFEI Publication #641, San Francisco Estuary Institute, Oakland, California.

Bendix, J. 1994. Among-site variation in riparian vegetation of the southern California transverse ranges. American Midland Naturalist 132: 136–151.

Bendix, J., and C. R. Hupp. 2000. Hydrological and geomorphological impacts on riparian plant communities. Hydrological Processes 14: 2,977–2,990.

Boughton, D. A., P. B. Adams, E. Anderson, C. Fusaro, E. Keller, E. Kelley, L. Lentsch, J. Nielsen, K. Perry, H. Regan, J. Smith, C. Swift, L. Thompson, and F. Watson. 2006. Steelhead of the south-central/southern California coast: population characterization for recovery planning. NOAA Technical Memorandum NMFS-SWFSC-394.

CDFW (California Department of Fish and Wildlife). 2019. California Natural Diversity Database. RareFind 5 [Internet], Version 5.1.1. [accessed: April 2019].

CDFW. 2020a. Special Vascular Plants, Bryophytes, and Lichens List. Accessed November 2020.

CDFW. 2020b. Sensitive Natural Communities List. Accessed November 2020.

CNPS (California Native Plant Society). 2020. A Manual of California Vegetation, online edition. <a href="http://www.cnps.org/cnps/vegetation/">http://www.cnps.org/cnps/vegetation/</a> [Accessed December 2020]. California Native Plant Society, Sacramento, California.

Daniel B. Stephens and Associates (DBS&A). 2020. Fillmore and Piru groundwater basins monitoring program and data gap analysis. Draft Technical Memorandum prepared for The Fillmore and Piru Basins Groundwater Sustainability Agency. September 2020.

DBS&A. 2021. Sustainable management criteria Climate Change 2070 Scenario. Presentation to The Fillmore and Piru Basins Groundwater Sustainability Agency. January 21 2021. https://s29420.pcdn.co/wp-content/uploads/2021/01/REVISION-to-Item-4C-2021-01-21-FPBGSA-SMC.pdf

DiTomaso, J. M. 1998. Biology and ecology of giant reed. Proceedings of the Arundo and Saltceder: The Deadly Duo Workshop, Ontario, California.

DiTomaso, J. M. and E. Healey. 2007. Weeds of California and Other Western States. University of California Press, Berkeley, California.

DWR (California Department of Water Resources). 2016. Bulletin 118 – Groundwater Basins and Subbasins, California.

DWR. 2020. Natural Communities Commonly Associated with Groundwater Dataset Viewer. <a href="https://gis.water.ca.gov/app/NCDatasetViewer/#">https://gis.water.ca.gov/app/NCDatasetViewer/#</a> [Accessed November 2020].

eBird. 2021. eBird: An online database of bird distribution and abundance. Website [accessed November 2020]. eBird, Cornell Lab of Ornithology, Ithaca, New York.

Edwards, R.D., D.F. Rabey, and R.W. Kover. 1970. Soil Survey of the Ventura Area, California. United States Department of Agriculture, Soil Conservation Service, in cooperation with the University of California Agricultural Experiment Station.

Giessow, J., J. Casanova, R. Leclerc, R. MacArthur, G. Fleming, and J. E. Giessow. 2011. *Arund o donax* (giant reed) Distribution and Impact Report. Prepared by the California Invasive Plant C ouncil (Cal-IPC) for the State Water Resources Control Board.

Harris, R. R. 1999. Defining reference conditions for restoration of riparian plant communities: examples from California, USA. Environmental Management 24: 55–63.

Herrera, A. M., and T. L. Dudley. 2003. Reduction of riparian arthropod abundance and diversity as a consequence of giant reed (*Arundo donax*) invasion. Biological Invasions 5: 167–177.

Howard, S. R. and Both, M. T. 2016. Range expansion of the Shimofuri goby (*Tridentiger bifasciatus*) in southern California, with emphasis on the Santa Clara River. California Fish and Game 102(2): 45-49.

Howard, S., M. Booth, and E. Lashly. 2015. Memo to Catherine McCalvin (United Water Conservation District). 4 August 2015.

Hunter, W. C., R. D. Ohmart, and B. W. Anderson. 1988. Use of the exotic saltcedar (*Tamarix chinensis*) by birds in arid riparian systems. Condor 90: 113–123.

Huntington, J., McGwire, K., Morton, C., Snyder, K., Peterson, S., Erickson, T., Niswonger, R., Carroll, R., Smith, G. and Allen, R., 2016. Assessing the role of climate and resource management on groundwater dependent ecosystem changes in arid environments with the Landsat archive. Remote sensing of Environment, 185, pp.186-197.

Hupp, C. R., and W. R. Osterkamp. 1996. Riparian vegetation and fluvial geomorphic processes. Geomorphology 14: 277–295.

Jepson Flora Project. 2020. Jepson eFlora. Website. <a href="http://ucjeps.berkeley.edu/eflora">http://ucjeps.berkeley.edu/eflora</a> [Accessed October 2020].

Kibler, C.L., Roberts, D.A., Singer, M.B. and Stella, J.C., 2019, December. Remote sensing of plant-groundwater relations in riparian forests during the 2011-2017 California drought. In AGU Fall Meeting Abstracts (Vol. 2019, pp. B53N-2588).

Kisner, D. A. 2004. The effect of giant reed (*Arundo donax*) on the southern California riparian bird community. Master's thesis. San Diego State University, San Diego, CA.

Klausmeyer, K., J. Howard, T. Keeler-Wolf, K. Davis-Fadtke, R. Hull, and A. Lyons. 2018. Mapping indicators of groundwater dependent ecosystems in California. https://groundwaterresourcehub.org/public/uploads/pdfs/iGDE\_data\_paper\_20180423.pdf

Kløve, B., Ala-Aho, P., Bertrand, G., Gurdak, J.J., Kupfersberger, H., Kværner, J., Muotka, T., Mykrä, H., Preda, E., Rossi, P. and Uvo, C.B., 2014. Climate change impacts on groundwater and dependent ecosystems. Journal of Hydrology, 518, pp.250-266.

Labinger, Z., and J. Greaves. 2001. Results of 2000 avian surveys and least Bell's vireo monitoring: restoration phase of the ARCO/Four Corners January 17, 1994 oil spill on the Santa Clara River, California. Report prepared for U.S. Fish and Wildlife Service, Ventura, California

- LARWQCB (Los Angeles Region Water Quality Control Plan). 2014. Basin Plan for the Coastal Watersheds of Los Angeles and Ventura Counties.
- Leenhouts, J. M., J.C. Stromberg, and R. L. Scott (editors). 2006. Hydrologic requirements of and consumptive ground-water use by riparian vegetation along the San Pedro River, Arizona. U.S. Geological Survey Scientific Investigations Reports 2005-5163.
- Lichvar, R. W., D. L. Banks, W. N. Kirchner, and N. C. Melvin. 2016. The National wetland plant list: 2016 wetland ratings. Phytoneuron 2016–30: 1–17.
- Lite, S. J. 2003. San Pedro River riparian vegetation across water availability and flood disturbance gradients. Ph.D. dissertation. Arizona State University, Phoenix.
- Lovich, J. C., and R. C. DeGouvenain. 1998. Saltcedar invasion in desert wetlands of the southwestern United States: Ecological and political implications. Pages 447-467 in S. K. Majumder, E. W. Miller, and S. J. Brenner, editors. Ecology of wetlands and associated systems. Pennsylvania Academy of Science, Easton, Pennsylvania.
- NMFS (National Marine Fisheries Services). 2005. Endangered and threatened species; designation of critical habitat for seven evolutionary significant units of Pacific salmon and steelhead in California. Federal Register 70: 52,488–52,627.
- NMFS. 2012. Southern California Steelhead Recovery Plan. Prepared by NMFS, Southwest Region, Protected Resources Division, Long Beach, California.
- Orr, B. K., Z. E. Diggory, G. C. Coffman, W. A. Sears, T. L. Dudley, and A. G. Merrill. 2011. Riparian vegetation classification and mapping: important tools for large-scale river corridor restoration in a semi-arid landscape. Pages 212–232 *in* J. Willoughby, B. Orr, K. Schierenbeck, and N. Jensen, editors. Proceedings of the CNPS Conservation Conference: Strategies and Solutions, 17–19 Jan 2009.
- Osterkamp, W. R. and C. R. Hupp. 2010. Fluvial processes and vegetation—glimpses of the past, the present, and perhaps the future. Geomorphology 116: 274–285.
- Rohde, M. M., B. Seapy, R. Rogers, X. Castañeda, editors. 2019. Critical Species LookBook: A compendium of California's threatened and endangered species for sustainable groundwater management. The Nature Conservancy, San Francisco, California.
- Rohde, M. M., S. Matsumoto, J. Howard, S. Liu, L. Riege, and E. J. Remson. 2018. Groundwater Dependent Ecosystems under the Sustainable Groundwater Management Act: Guidance for Preparing Groundwater Sustainability Plans. The Nature Conservancy, San Francisco, California.
- Santa Clara River Trustee Council. 2008. Santa Clara River Watershed Amphibian and Macroinvertebrate Bioassessment Project. Prepared by The Wishtoyo Foundation in association with South Coast Wildlands.
- Shafroth, P. B., G. T. Auble, J. C. Stromberg, and D. T. Patten. 1998. Establishment of woody riparian vegetation in relation to annual patterns of streamflow, Bill Williams River, Arizona. Wetlands 18: 577–590.

Shafroth, P. B., J. R. Cleverly, T. L. Dudley, J. P. Taylor, C. van Riper III, E. P. Weeks, and J. N. Stuart. 2005. Control of Tamarix in the western United States: Implications for water salvage, wildlife use, and riparian restoration. Environmental Management 35: 231–246.

SoCalGas (Southern California Gas Company). 2020. Coastal Region Conservation Program multi-species Habitat Conservation Plan.

Stillwater Sciences. 2007a. Santa Clara River Parkway Floodplain Restoration Feasibility Study: analysis of riparian vegetation dynamics for the lower Santa Clara River and major tributaries,

Swain, D.L., Langenbrunner, B., Neelin, J.D. and Hall, A., 2018. Increasing precipitation volatility in twenty-first-century California. Nature Climate Change, 8(5), pp.427-433.

Ventura County, California. Prepared by Stillwater Sciences for the California State Coastal Conservancy.

Stillwater Sciences. 2007b. Santa Clara River Parkway. Digital vegetation map managed through the California Vegetation Classification and Mapping Program (VegCAMP), prepared for the California Coastal Conservancy. Accessed January 2016.

Stillwater Sciences. 2007c. Focal Species Analysis and Habitat Characterization for the Lower Santa Clara River and Major Tributaries, Ventura County, California. Santa Clara River Parkway Floodplain Restoration Feasibility Study. Prepared by Stillwater Sciences for the California State Coastal Conservancy and the Santa Clara River Trustee Council. SCE (Southern California Edison Gas Company). 2020. Coastal Region Conservation Program Multiple Species Habitat Conservation Plan.

Stillwater Sciences. 2007d. Santa Clara River Parkway Floodplain Restoration Feasibility Study: Assessment of Geomorphic Processes for the Santa Clara River Watershed, Ventura and Los Angeles Counties, California. Prepared by Stillwater Sciences for the California State Coastal Conservancy.

Stillwater Sciences. 2011. Geomorphic assessment of the Santa Clara River watershed: synthesis of the lower and upper watershed studies, Ventura and Los Angeles counties, California. Prepared by Stillwater Sciences, Berkeley, California for Ventura County Watershed Protection District, Los Angeles County Department of Public Works, and the U.S. Army Corps of Engineers—L.A. District.

Stillwater Sciences. 2016. United Water Conservation District multiple species habitat conservation plan study: effects of Freeman Diversion on habitat conditions in the Santa Clara River Estuary, draft technical report. Prepared by Stillwater Sciences, Berkeley, California for United Water Conservation District, Santa Paula, California.

Stillwater Sciences. 2019. Vegetation Mapping of the Santa Clara River, Ventura County and Los Angeles County, California. Technical Memorandum. Prepared by Stillwater Sciences, Berkeley, California for the Western Foundation of Vertebrate Zoology, Camarillo, California.

Stoecker, M., and E. Kelley. 2005. Santa Clara River steelhead trout: assessment and recovery opportunities. Prepared for The Nature Conservancy and The Santa Clara River Trustee Council, Santa Barbara, California.

Stromberg, J. C., R. Tiller, and B. Richter. 1996. Effects of groundwater decline on riparian vegetation of semiarid regions: the San Pedro, Arizona. Ecological Applications 6: 113–131.

Swift, C. C., T. R. Haglund, M. Ruiz, R. M. Fisher. 1993. The Status and Distribution of the Freshwater Fishes of Southern California. Bulletin of the Southern California Academy of Sciences 92: 101–167.

TNC (The Nature Conservancy). 2006. Santa Clara River Upper Watershed Conservation Plan.

TNC. 2008. Conservation Plan for the Lower Santa Clara River Watersheds and Surrounding Areas.

TNC. 2020. Freshwater species list for Fillmore and Piru Groundwater Basins. <a href="https://groundwaterresourcehub.org/sgma-tools/environmental-surface-water-beneficiaries">https://groundwaterresourcehub.org/sgma-tools/environmental-surface-water-beneficiaries</a>. Accessed December 2020.

US Department of Agriculture - Forest Service. 2014. Existing Vegetation South Coast, 2002–2010, v2 [ESRI File Geodatabase]. McClellan, CA. Accessed June 2016.

USFS (U.S. Forest Service). 2011. FSM 2600 – Wildlife, Fish, and Sensitive Plant Habitat Management, Chapter 2670 – Threatened, Endangered, and Sensitive Plants and Animals. Forest Service Manual Rocky Mountain Region (Region 2). Denver, Colorado.

USFWS (U.S. Fish and Wildlife Service). 1976. Determination of Critical Habitat for American Crocodile, California Condor, Indiana Bat, and Florida Manatee. Federal Register 41: 41,914–41,916.

USFWS. 1977. Final Rule; Correction and Augmentation of Published Rulemaking. Federal Register 42: 47,840–47,845.

USFWS. 1994. Designation of Critical Habitat for Least Bell's Vireo: Final Rule. Federal Register 59: 4,845–4,867.

USFWS. 1999. Arroyo southwestern toad (*Bufo microscaphus californicus*) recovery plan. U.S. Fish and Wildlife Service, Portland, Oregon.

USFWS. 2013. Designation of Critical Habitat for Southwestern Willow Flycatcher. Federal Register 78: 344–534.

USFWS. 2017. Recovery Plan for the Santa Ana sucker (*Catostomus santaanae*). U.S. Fish and Wildlife Service, Pacific Southwest Region, Sacramento, California.

USFWS. 2018. National Wetlands Inventory website. U.S. Department of the Interior, Fish and Wildlife Service, Washington, D.C. <a href="http://www.fws.gov/wetlands/">http://www.fws.gov/wetlands/</a>

USGS (United States Geological Survey). 2018. 2018 USGS LiDAR: Southern CA Wildfires Point Cloud files with Orthometric Vertical Datum North American Vertical Datum of 1988 (NAVD88) using Geoid18. Accessed December 2020.

UWCD (United Water Conservation District). 2016. 2014 and 2015 Piru and Fillmore Basins Biennial Groundwater Conditions Report. United Water Conservation District Open-File Report 2016-01.

UWCD. 2018. United Water Conservation District Multiple Species Habitat Conservation Plan (Administrative Draft).

UWCD. 2020. Water Level Elevation contours, GIS data. Provided by Daniel B. Stephens & Associates August 2020.

VCWPD (Ventura County Watershed Protection District) and LADPW (Los Angeles County Department of Public Works). 2005. Santa Clara River Enhancement and Management Plan. Prepared by AMEC Earth and Environmental.

WFVZ (The Western Foundation of Vertebrate Zoology). 2020a. 2020 Report on Nesting Bird Surveys and Nest Management during Restoration on the Cienaga Springs Ecological Reserve, Ventura County, California. Prepared for U.C Santa Barbara, the Santa Clara River Conservancy, and the California Department of Fish and Wildlife.

WFVZ. 2020b. 2020 Compliance Report Pertaining to Nesting Bird Surveys and Nest Management During Restoration on the Santa Clara River, Ventura County, California. Prepared for The Nature Conservancy.

WFVZ. 2020c. Final Report on Bird Counts on Nature Conservancy Properties along the Santa Clara River, Ventura CO, California. Prepared for The Nature Conservancy.

Zembal, R. 1990. Riparian habitat and breeding birds along the Santa Margarita and Santa Ana Rivers of southern California. Pages 98-114 in A. A. Schoenherr, editor. Endangered plant communities of southern California. Southern California Botanists, Special Publ. No. 3.

# **Appendices**

### Appendix A

Vegetation Communities in the Fillmore and Piru Groundwater Basins

**Table A-1**. All vegetation communities mapped in the Fillmore and Piru groundwater basins and corresponding vegetation classification crosswalk.

		Acreage			
CALVEG Classification	Manual of California Vegetation (MCV) <sup>1</sup>	Fillmore Groundwater Basin	Piru Groundwater Basin		
Alkaline mixed grasses and forbs Alliance	Cressa truxillensis - Distichlis spicata Herbaceous Alliance	-	2.2		
Annual grasses and forbs Alliance	Brassica nigra - Raphanus spp. Herbaceous Semi- Natural Alliance	1,291.1	377.4		
Arrowweed Alliance	Pluchea sericea Shrubland Alliance	=	31.5		
Baccharis (riparian) Alliance	Baccharis salicifolia Shrubland Alliance	1,216.1	952.2		
Долгон	Riverwash	138.5	84.9		
Barren	Riverwash herbaceous	138.3	84.9		
Black cottonwood Alliance	Populus trichocarpa Forest Alliance	320.6	-		
Black walnut Alliance	Juglans californica Woodland Alliance	6.8	3.7		
Big sagebrush Alliance	Artemisia tridentata Shrubland Alliance	4.0	54.1		
Blue oak Alliance	n/a	4.0	-		
Buckwheat	Encelia californica - Eriogonum cinereum Shrubland Alliance	113.8	27.3		
	Eriogonum fasciculatum Shrubland Alliance				
California sagebrush Alliance	Artemisia californica Shrubland Alliance	1,727.4	639.8		
Camornia sageorasii rimanee	Salvia apiana Shrubland Alliance	1,727.1			
California sycamore Alliance	Platanus racemosa Woodland Alliance	-	4.6		
Ceanothus chaparral Alliance	n/a	44.2	-		
Chamise Alliance	n/a	2.8	-		
Coast live oak Alliance	Quercus agrifolia Woodland Alliance	815.8	65.4		
Coastal mixed hardwood Alliance	n/a	77.8	4.0		
Coyote brush Alliance	Baccharis pilularis Shrubland Alliance	4.1	0.9		
Eucalyptus Alliance	Eucalyptus spp Ailanthus altissima - Robinia pseudoacacia Woodland Semi-Natural Alliance	74.7	14.4		
Fremont cottonwood Alliance	Populus fremontii Forest Alliance	0.7	244.8		
Giant reed/pampas grass Alliance	Phragmites australis - Arundo donax Herbaceous Semi-Natural Alliance	271.6	183.0		
Lower montane mixed chaparral Alliance	n/a	170.7	3.1		
Manzanita chaparral Alliance	n/a	< 0.1	-		
Non-native/ornamental conifer/hardwood Alliance	n/a	13.0	2.6		
Non-native/ornamental grass Alliance	Non-native Grass and Forb Mapping Unit	195.0	90.2		
Non-native/ornamental hardwood Alliance	Schinus (molle, terebinthifolius) - Myoporum laetum Woodland Semi-Natural Alliance	75.7	20.2		
Perennial grasses and forbs Alliance	Corethrogyne filaginifolia - Eriogonum (elongatum, nudum) Herbaceous Alliance  Leymus cinereus - Leymus triticoides Herbaceous Alliance	6.3	-		
Riparian mixed hardwood Alliance	Sambucus nigra Shrubland Alliance Salix laevigata Woodland Alliance	396.2	286.4		

		Acr	Acreage		
CALVEG Classification	CALVEG Classification Manual of California Vegetation (MCV) <sup>1</sup>				
Riparian mixed shrub Alliance	Heterotheca (oregona, sessiliflora) Herbaceous Alliance	152.2	557.1		
	Salix exigua Shrubland Alliance				
Riversidean alluvial scrub Alliance	n/a	52.5	3.6		
Saltbrush Alliance	Atriplex lentiformis Shrubland Alliance	54.1	58.1		
Scalebroom Alliance	Lepidospartum squamatum Shrubland Alliance	320.1	118.0		
Scrub oak Alliance	n/a	1.1	-		
Soft scrub-mixed chaparral Alliance	n/a	62.9	-		
Sumac shrub Alliance	n/a	522.4	0.9		
Tamarisk Alliance	Tamarix spp. Shrubland Semi-Natural Alliance	-	37.6		
Tule-cattail Alliance	Schoenoplectus (acutus, californicus) Herbaceous Alliance	8.1	3.3		
Tule-cauaii Amance	Typha (angustifolia, domingensis, latifolia) Herbaceous Alliance	0.1	3.3		
Wet meadow	n/a	0.4	-		
Willow / Willow (shrub) Alliance	Salix lasiolepis Shrubland Alliance	63.5	3.9		
winow / winow (siliub) Alliance	Salix lucida Woodland Alliance	05.5	3.9		
	Olea europaea Woodland Semi-Natural Alliance [Provisional]	-	2.7		
No corresponding CalVeg type <sup>2</sup>	Pseudognaphalium leucocephalum Herbaceous Alliance [Provisional]	-	0.2		
	Ricinus communis Shrubland Semi-Natural Alliance [Provisional]	-	2.2		
All agriculture		12,436.9	6,123.8		
All water		6.7	14.6		
All development		1,968.6	903.2		
Total		22,620.3	10,922.0		

An n/a in this column signifies that no corresponding MCV type was mapped in the Vegetation Mapping of Santa Clara River dataset (Stillwater Sciences 2019).
 These are provisional MCV alliances and as such do not have a corresponding CalVeg alliance.

### Appendix B

Special-status Terrestrial and Aquatic Wildlife Species from Database Queries with No Reliance on Fillmore or Piru Groundwater Dependent Ecosystem Units

Table. B-1. Special-status terrestrial and aquatic wildlife species from database queries that are not groundwater dependent and/or unlikely to occur in the Fillmore and Piru groundwater dependent ecosystem units.

Common name	Status <sup>1</sup>	Potential to occur in		d occurrence	Query	GDE.	Habitat and documented occurrences in
Scientific name	Federal/State	GDE Basins <sup>2</sup>	Fillmore GDE units	Piru GDE units	source <sup>3</sup>	association <sup>4</sup>	GDE Management Units
Amphibian							
California red- legged frog Rana draytonii	FT/SSC	Unlikely	No documented occurrences	No documented occurrences	CAFSD	Direct	Breeds in still or slow-moving water with emergent and overhanging vegetation, including wetlands, wet meadows, ponds, lakes, and low-gradient, slow moving stream reaches with permanent pools; uses adjacent uplands for dispersal and summer retreat. Relies on surface water that may be supported by groundwater (Rohde et al. 2019).
Foothill yellow- legged frog <i>Rana boylii</i>	FSS, BLMS/SE	None	Extirpated	Extirpated	CNDDB, CAFSD	Direct	Shallow tributaries and mainstems of perennial streams and rivers, typically associated with cobble or boulder substrate; occasionally found in isolated pools, vegetated backwaters, and deep, shaded, spring-fed pools. The frog is reliant on surface water that may be fed by groundwater. Population has been extirpated from the Santa Clara River Valley Basin (CDFW 2019).
Western spadefoot Spea hammondii	BLMS/SSC	Unlikely	No documented occurrences	No documented occurrences	CAFSD	No known reliance on groundwater	Areas with sparse vegetation and/or short grasses in sandy or gravelly soils; primarily in washes, river floodplains, alluvial fans, playas, alkali flats, among grasslands, chaparral, or pine-oak woodlands; breeds in ephemeral rain pools with no predators.

Common name	Status <sup>1</sup>	Potential to occur in		d occurrence	Query	GDE.	Habitat and documented occurrences in
Scientific name	Federal/State	GDE Basins <sup>2</sup>	Fillmore GDE units	Piru GDE units	source <sup>3</sup>	association <sup>4</sup>	GDE Management Units
Reptile							
California legless lizard Anniella sp.	-/SSC	Likely	Santa Clara River Riparian Shrubland	Tributary Riparian	CNDDB	No known reliance on groundwater	Occurs in moist warm loose soil with plant cover and in sparsely vegetated areas of chaparral, pine-oak woodlands, desert scrub, and stream terraces with sycamores, cottonwoods, or oaks. Forages in loose soil, sand, and leaf litter for larval insects, beetles, termites, and spiders. Historical observation in the vicinity of Sespe Creek and Santa Clara River confluence in 1981 (CDFW2019). Observations in the vicinity of Tributary Riparian GDE Unit include Hopper Canyon in 2008 (CDFW 2019).
Coast horned lizard Phrynosoma blainvillii	FSS, BLMS/SSC	Likely	East Grove Riparian Complex	Santa Clara River Riparian Shrubland	CNDDB	No known reliance on groundwater	Open areas with sandy soil and/or patches of loose soil and low/scattered vegetation in scrublands, grasslands, conifer forests, and woodlands; frequently found near ant hills. Feeds on ants and other small invertebrates (e.g., spiders, beetles, and grasshoppers).
Coast patch-nosed snake Salvadora hexalepis virgultea	-/SSC	Likely	No documented occurrences	No documented occurrence	CNDDB	No known reliance on groundwater	Coastal chaparral, desert scrub, washes, sandy flats and rocky areas. Predominately preys upon lizards. Documented outside of groundwater basins on Hopper Creek, two miles northwest of Piru (CDFW 2019).
San Diegan Coastal whiptail Aspidoscelis tigris stejnegeri	-/SSC	Likely	No documented occurrences	Piru Creek Riparian	CNDDB	No known reliance on groundwater	Habitat generalists found in desert, woodland, and riparian communities. Feeds on small invertebrates (e.g., spiders, scorpions, centipedes, and termites) and small lizards. Documented on Piru Creek in 2009 (CDFW 2019)

Common name	Status <sup>1</sup>	Potential to occur in		d occurrence	Query	GDE.	Habitat and documented occurrences in
Scientific name	Federal/State	GDE Basins <sup>2</sup>	Fillmore GDE units	Piru GDE units	source <sup>3</sup>	association <sup>4</sup>	GDE Management Units
Bird							
Bank swallow Riparia riparia	BLMS/ST	None	Extirpated	Extirpated	CNDDB, CAFSD	Indirect	Nests in vertical bluffs or banks, usually adjacent to water (i.e., rivers, streams, ocean coasts, and reservoirs), where the soil consists of sand or sandy loam. Feeds on caterpillars, insects, frog/lizards, and fruit/berries. Relies on surface water that may be supported by groundwater (Rohde et al. 2019). Historical population documented in the 1920's Santa Clara River is extirpated (CDFW 2019).
Black swift Cypseloides niger	FSS/SSC	Unlikely	No documented occurrences	No documented occurrences	CAFSD	No known reliance on groundwater	Nests in moist crevices behind or beside permanent or semipermanent waterfalls in deep canyons, on perpendicular sea cliffs above surf, and in sea caves; forages widely for insects over many habitats.
Burrowing owl Athene cunicularia	FSS/SSC	Likely	Santa Clara River Riparian Shrubland	Santa Clara River Riparian Shrubland	CNDDB	No known reliance on groundwater	Level, open, dry, heavily grazed or low-stature grassland or desert vegetation with available burrows. Preys on invertebrates and vertebrates. Occurrences along or near the bank of the Santa Clara River near Fillmore and one mile south of Buckhorn (CDFW 2019).

Common name Scientific name	Status <sup>1</sup> Federal/State	Potential to occur in GDE Basins <sup>2</sup>		d occurrence tion Piru GDE units	Query source <sup>3</sup>	GDE . association <sup>4</sup>	Habitat and documented occurrences in GDE Management Units
Mammal							
Pallid bat Antrozous pallidus	FSS, BLMS/SSC	Likely	No documented occurrences	No documented occurrences	CNDDB	No known reliance on groundwater	Roosts in rock crevices, tree hollows, mines, caves, and a variety of vacant and occupied buildings; feeds in a variety of open woodland habitats. Habitat and prey (e.g., insects and arachnids) not associated with aquatic ecosystems. Commonly found roosting under the bark of dead riparian trees in the Santa Clara River Watershed (UWCD 2018). Historical observations in the vicinity of Fillmore documented in 1906 and 1942 (CDFW 2019).

<sup>&</sup>lt;sup>1</sup> Status codes:

Federal State

FT = Listed as threatened under the federal Endangered Species Act
FSS = Forest Service Sensitive Species

SE = Listed as Endangered under the California Endangered Species Act
SE = Listed as Threatened under the California Endangered Species Act

BLMS = Bureau of Land Management Sensitive Species SSC = CDFW species of special concern

#### Potential to Occur:

Likely: the species has documented occurrences and the habitat is high quality or quantity Possible: no documented occurrences and the species' required habitat is moderate to high quality or quantity Unlikely: no documented occurrences and the species' required habitat is of low to moderate quality or quantity None: no potential to occur due to lack of habitat and/or the population is assumed extirpated

<sup>3</sup> Ouery source:

CAFSD: California Freshwater Species Database (TNC 2020) CNDDB: California Natural Diversity Database (CDFW 2019) eBird: (eBird 2021)

<sup>4</sup> Groundwater Dependent Ecosystem (GDE) association:

Direct: Species directly dependent on groundwater for some or all water needs

Indirect: Species dependent upon other species that rely on groundwater for some or all water needs

## Appendix C

### **Rooting Depths for Selected Species**

Table C-1. Maximum rooting depth of dominant species.

Dominant species	Vegetation type (MCV)	Vegetation type (CalVeg)	GDE?	Maximum rooting depth (ft)	Data source
Adenostoma fasciculatum		Chamise	no	25.0	Hellmers et al. 1955 as cited in Fan et al. 2017
Artemisia tridentata	Artemisia tridentata Shrubland Alliance		yes	9.8	Link et al. 1995 as cited in Tumber- Davila 2017
Arundo donax	Phragmites australis - Arundo donax Herbaceous Semi-Natural Alliance	Giant reed/pampas grass	yes	16.1	Stromberg 2013
Baccharis pilularis	Baccharis pilularis Shrubland Alliance	Coyote brush	no	12.1	Naumovich 2017
Baccharis salicifolia	Baccharis salicifolia Shrubland Alliance	Baccharis (riparian)	yes	2.0	Stromberg 2013
Ceanothus crassifolius		Ceanothus chaparral	no	4.5	Hellmers et al. 1955 as cited in Fan et al. 2017
Elymus triticoides	Leymus cinereus - Leymus triticoides Herbaceous Alliance		yes	3.8	Weaver 1919 as cited in Fan et al. 2017
Eriogonum fasciculatum	Eriogonum fasciculatum Shrubland Alliance	Buckwheat	no	4.0	Hellmers et al. 1955 as cited in Fan et al. 2017
Eucalyptus globulus and other Eucalyptus species	Eucalyptus spp Ailanthus altissima - Robinia pseudoacacia Woodland Semi- Natural Alliance	Eucalyptus	yes	16.4	Dawson and Pate 1996 as cited in Fan et al. 2017
Juglans californica	Juglans californica Woodland Alliance	Black walnut	no	5.9	Faber 2017
Pluchea sericea	Pluchea sericea Shrubland Alliance		yes	4.3	Stromberg 2013
Populus fremontii	Populus fremontii Forest Alliance	Freemont cottonwood	yes	6.9	Stromberg 2013
Populus trichocarpa	Populus trichocarpa Forest Alliance		yes	4.1	Zhang et al. 1999 as cited in Fan et al. 2017
Quercus agrifolia	Quercus agrifolia Woodland Alliance	Coast live oak / Coastal mixed hardwood alliance	yes	35.1	Schenk and Jackson 2002
Quercus douglasii		Blue oak	nc =	80.0	Schenk and Jackson 2002

Dominant species	Vegetation type (MCV)	Vegetation type (CalVeg)	GDE?	Maximum rooting depth (ft)	Data source
Salix exigua	Salix exigua Shrubland Alliance	Willow (shrub)	yes	6.9 <sup>1</sup>	Pulling 1918 as cited in Fan et al. 2017
Salix laevigata	Salix laevigata Woodland Alliance		yes	6.9 <sup>1</sup>	Pulling 1918 as cited in Fan et al. 2017
Salix lasiolepis	Salix lasiolepis Shrubland Alliance		yes	6.9 <sup>1</sup>	Pulling 1918 as cited in Fan et al. 2017
Salix lucida	Salix lucida Woodland Alliance		yes	6.9 <sup>1</sup>	Pulling 1918 as cited in Fan et al. 2017
Salix spp		Riparian mixed shrub / Willow	yes	6.9	Pulling 1918 as cited in Fan et al. 2017
Salvia apiana	Salvia apiana Shrubland Alliance		no	5.0	Hellmers et al. 1955 as cited in Fan et al. 2017
Schoenoplectus spp	Schoenoplectus (acutus, californicus) Herbaceous Alliance		yes	2.11	Stromberg 2013
Tamarix spp	Tamarix spp. Shrubland Semi-Natural Alliance		yes	16.1	Stromberg 2013
Typha spp	Typha (angustifolia, domingensis, latifolia) Herbaceous Alliance		yes	0.81	Shaver and Billlings 1975 as cited in Fan et al. 2017

<sup>&</sup>lt;sup>1</sup> Rooting depth assigned by genus or close species association.

Table C-2. Relative elevation of dominant species. Data from Stillwater Sciences (2007).

		Relative Elevation (ft)					
Dominant Species	Alliance	mean	SE	min	max		
Adenostoma fasciculatum							
Artemisia tridentata	Artemisia tridentata Shrubland Alliance	5.2	0.5	1.0	11.0		
Arundo donax	Arundo donax Semi-Natural Alliance	7.6	0.3	0.0	32.8		
Baccharis pilularis	Baccharis pilularis Shrubland Alliance	15.9	1.2	2.5	26.7		
Baccharis salicifolia	Baccharis salicifolia Shrubland Alliance	9.6	1.8	2.3	24.6		
Eriogonum fasciculatum and Artemisia californica	Artemisia californica- Eriogonum fasciculatum Shrubland Alliance	18.2	2.2	7.7	27.4		
Eucalyptus globulus and other Eucalyptus species	Eucalyptus spp Ailanthus altissima - Robinia pseudoacacia Woodland Semi-Natural Alliance	15.4	2.6	12.5	23.0		
Populus fremontii	Populus fremontii Forest Alliance	9.7	0.8	0.0	25.0		
Populus trichocarpa	Populus balsamifera ssp trichocarpa Forest Alliance	7.4	0.4	2.1	20.5		
Salix exigua	Salix exigua Shrubland Alliance	4.8	0.4	1.0	9.4		
Salix laevigata	Salix laevigata Woodland Alliance	5.6	0.5	0.0	20.6		
Salix lasiolepis	Salix lasiolepis Shrubland Alliance	12.4	0.7	0.6	24.8		
Salix lucida	Salix lucida Woodland Alliance	4.9	0.8	0.2	12.3		
Tamarix spp	Tamarix spp. Semi-Natural Alliance	6.3	1.1	4.0	9.0		

#### Literature Cited

Faber, B. 2017. TNC Crowdsourcing Campaign Survey Response.

Fan, Y., G. Miguez-Macho, E. G. Jobbágy, R. B. Jackson and C. Otero-Casal. 2017. Hydrologic regulation of plant rooting depth. PNAS October 3, 2017. 114: 10,572–10,577

Naumovich, L. 2017. TNC Crowdsourcing Campaign Survey Response.

Schenk, H. J. and R. B. Jackson. 2002. The Global Biogeography of Roots. Ecological Monographs, 72: 311–328. doi:10.1890/0012-9615(2002)072[0311:TGBOR]2.0.CO;2.

Stillwater Sciences. 2007. Santa Clara River Parkway Floodplain Restoration Feasibility Study: analysis of riparian vegetation dynamics for the lower Santa Clara River and major tributaries, Ventura County, California. Prepared by Stillwater Sciences for the California State Coastal Conservancy.

Stromberg, J. 2013. Root patterns and hydrogeomorphic niches of riparian plants in the America Southwest. Journal of Arid Environments 94 (2013) 1–9. Appendix A. Rooting data for herbaceous plants.

Tumber-Davila, S. J. .2017. Download to TNC of California-Specific Rooting Depth Data from the 2017 version of the Schenk, H.J. and R.B. Jackson (2002) The Global Biogeography of Roots. Ecological Monographs 72: 311–328.